

Suitability of Stimuli evoked Hypersensitivities in Hind paws as a Clinically Relevant Pain Behavioural Measure in Rat Model of Walker 256 Breast Cancer Cell-induced Bone Pain: An Overview

Priyank A. Shenoy*

Faculty of Pharmacy and Pharmaceutical Sciences, Monash University, Melbourne, Victoria, Australia

Corresponding author:

Dr. Priyank A. Shenoy

✉ priyank.shenoy@monash.edu

Faculty of Pharmacy and Pharmaceutical Sciences, Monash University, Melbourne, Victoria, Australia.

Tel: +61 3 9905 4000

Citation: Shenoy PA (2018) Suitability of Stimuli evoked Hypersensitivities in Hind paws as a Clinically Relevant Pain Behavioural Measure in Rat Model of Walker 256 Breast Cancer Cell-induced Bone Pain: An Overview. *Am J Pharmacol Pharmacother.* Vol.5 No.1:4.

Abstract

Metastases of breast cancer cells to the axial skeleton causes excruciating pain. The nature of pain hypersensitivities in metastasized bones is very complex due to the interplay of inflammatory, neuropathic and cancer-specific pain components. The existing drugs typically used to treat breast cancer-induced bone pain are inefficacious and often exhibit severe side effects. Hence, it remains to be an important goal of the ongoing research activities to seek novel analgesic compounds with better efficacy and tolerability. One of the key aspects in the process of understanding the mechanisms of pain progression and drug discovery to mitigate the hypersensitivities is employment of suitable preclinical animal models that mimic the complex human pathophysiology of breast cancer induced bone pain. Walker 256 breast cancer cell-induced bone pain model in rats is one such model that is known to show key resemblances to the clinical pain associated with bony metastases. The commonly used methods to assess pain hypersensitivities in Walker 256 cell-induced bone pain model in rats include stimuli evoked techniques like von Frey assessment and Randall-Selitto test. While other methods like assessment of gait parameters or spontaneous pain can also be used as beneficial complementary tools, this short review majorly sheds light on suitability of the stimuli evoked pain assessment methods in the hind paws of rats as being clinically relevant measures of assessing breast cancer induced bone pain in Walker 256 cell induced bony metastases.

Keywords: Walker 256; Breast cancer-induced bone pain

Received: August 22, 2018; **Accepted:** September 04, 2018; **Published:** September 06, 2018

Introduction

Pain is a significant medical problem that co-exists with several diseases including various types of cancer [1]. Breast cancer cells metastasize from the tissue of origin and establish themselves in distant parts of the axial skeleton [2]. The cancer cells growing in the bone microenvironment cause osteolysis and sensitisation of the peripheral nerve endings innervating the bones, thereby causing excruciating pain [3]. Breast cancer-induced bone pain (BCIBP) causes severe morbidity because of the heterogeneous combination of inflammatory, neuropathic and cancer-specific components [4]. The existing analgesic/adjvant medications are often insufficiently efficacious to combat this pain condition [5-

8]. Thus it is very important to develop and characterize suitable preclinical models of BCIBP so as to assist in drug discovery programs aimed at identifying novel compounds having potential to mitigate this often intractable pain condition. The rat model of Walker 256 breast cancer cell induced bone pain is a highly useful preclinical tool for assessment of mechanisms of BCIBP and for seeking novel analgesics in the treatments thereof, as it mimics key aspects of the human pathophysiology of this condition [8-11].

Von Frey test using a series of filaments corresponding to different levels of forces, and paw pressure test also called as Randall-Selitto testing using increasing force delivered via a blunt cone

shaped pusher, are two of the most common behavioural tests employed in assessment of Walker 256 cell induced BCIBP in rats [9]. The plantar hind paw of rats, the anatomical region where von Frey and Randall-Selitto stimuli are applied, is mainly innervated by the tibial nerve and hence tibial bone pain sensations manifest as hypersensitivities in the plantar aspect of the hind paws [12-15]. Hence, the traditional and most commonly used method to measure tibial bone pain is assessment in the paws, with hundreds of studies prevailing in the literature using this protocol. By using intra-tibial injections of complete Freund's adjuvant in the tibiae of female Wistar rats, a study has elegantly established that the bone pain induced by activation of tibial nerves directly manifests as hind paw skin hypersensitivity [15]. Cutaneous tests like the von Frey testing and Randall-Selitto testing detect the current levels of pain and have high clinical relevance as used in humans [16-21].

Discussion

A study involving the Department of Medicine of the University of Florida (Gainesville, USA) has validated that mechanically evoked pain is a highly relevant measure of the clinical pain intensity in patients with deep pain of muco-skeletal origin [22]. Similarly, a study conducted in Edinburgh Cancer Centre (Edinburgh, UK) also validated that assessing mechanical allodynia using von Frey filaments is a direct measure of cancer induced bone pain in humans [23]. Hence assessment of the stimuli evoked hypersensitivities in the hind paws is physiologically relevant assessment of bone pain. As per the previous studies published in journals like PAIN [24-26], The Journal of Pain [27,28], European Journal of Pain [29-31], Pain Medicine [32], Molecular Pain [33,34], Nature Neuroscience [35] and others, it is a traditional practice within the pain research fraternity to test the pain hypersensitivities in the paw, following inoculation of cancer cells in the tibia, without deploying other measures like gait or weight bearing parameters. Along these lines, a recent report suggested that a vast majority of around ~90% of the cancer induced bone pain studies in the literature using MRMT-1 cells in rats used the response evoked by the cutaneous stimuli applied to the foot as a measure of bone pain [36]. From the vast literature available on Walker 256 cell induced bone pain model in rats, inoculation of Walker 256 cell in the tibia always manifests as hypersensitivities in the hind paws, without any discordance in paw-tibia correlation being reported [8]. The majority of studies in the literature that used the Walker 256 breast cancer cell-induced bone pain model in rats, used the hind paw as a location to test hypersensitivity to evoked pain such as that induced by the von Frey test, rather than spontaneous or movement evoked pain [18-45]. There are many different studies very recently published, that used Walker 256 cells to induce bone pain in rats that only used stimuli evoked behavioural measures such as von Frey paw withdrawal thresholds in the hind paws, but not spontaneous movement evoked or weight bearing measures to assess pain hypersensitivities [35-60]. The vast experience of different laboratories conducting pre-clinical cancer-pain research around the world with the Walker 256 cell induced BCIBP model in rats strongly emphasizes on the suitability of stimuli evoked hypersensitivities in paws as the correct measure of bone pain in this particular model.

However, it is noteworthy that dissociation is observed in between skeletal pain behaviors and skin hypersensitivity in a male C3H mouse model of intra-femoral injection of NCTC 2472 osteosarcoma cells [61]. It is known that different types of cell lines or tumors exhibit distinct pain behavioral patterns [62]. It is the unique interaction between each of the cancers colonising the bone and the nerve innervation that predominantly decides the nature of pain manifestation [63]. On these grounds, a previous study highlighted the fact that neither spontaneous pain nor ambulatory pain is the best measure of cancer induced bone pain for all models triggered by different cancer cell lines in general [62]. It showed that intra-osseous injection of B16-F10 melanoma cell line in the femur did not produce either the spontaneous pain or the ambulatory pain. The bone pain induced by B16-F10 cell line manifested only as hind paw skin hypersensitivity. Similarly, C26 colon cancer cell line did not produce spontaneous pain behavior. Hence, the spontaneous or ambulatory pain are not the universal measures of bone pain at least in some models like B16 cell model and C26 cell model. In alignment to this pre-clinical animal based study, a clinical study of cancer induced bone pain also reported that in patients with breakthrough pain, which is commonly triggered by a stimulus [64], patients were not more likely to experience pain at the weight-bearing bone sites, compared to patients without pain [65]. Additionally, allodynia and gait behaviours are two independent phenomena. Neuropathic pain is one of the key components of cancer induced bone pain [4], and allodynia (measured by tests like von Frey) is a more reliable measure of the neuropathic pain component, rather than gait behaviours (weight bearing or spontaneous pain during ambulation) [66]. The changes in gait parameters can typically be due to the tendency of animals to avoid allodynia produced by contact of the paw with the floor [67]. The gait changes might not necessarily relate well to pain hypersensitivities [66-72]. There are several evidences that suggest that changes in gait parameters like guarding the hind paw during ambulation or changes in weight bearing are significantly driven by the adaptive changes and psychological influences (pain-avoidance and fear due to cognition), rather than the current levels of pain intensity [66-82]. Whereas, both the von Frey and Randall-Selitto tests that are also used in humans to assess pain hypersensitivities, detect the current levels of pain and have high clinical relevance [16-21]. This is probably one of the most important reasons why large number of published studies used stimuli-evoked methods like the von Frey and Randall-Selitto tests to assess pain hypersensitivities in the hind paws of animals following unilateral tibial inoculation of cancer cells.

Conclusion

The vast literature on Walker 256 cell induced BCIBP model in rats strongly suggests that stimuli evoked pain behaviours in the hind paws of rats is an appropriate measure of cancer induced bone pain in this particular model. However, a complementary assessment of measures like ambulatory pain, spontaneous pain or pain evoked by weight bearing on the hind paws of rats might add more value to the studies in future as these tests might be considered relatable to pain assessments in humans.

References

- Basbaum AI, Bautista DM, Scherrer G (2009) Cellular and molecular mechanisms of pain. *Cell* 139: 267-284.
- Cleeland C, Von Moos R, Walker MS (2016) Burden of symptoms associated with development of metastatic bone disease in patients with breast cancer. *Support Care Cancer* 24: 3557-3565.
- Nencini S, Ivanusic JJ (2016) The physiology of bone pain. How much do we really know? *Front Physiol* 7.
- Cao F, Gao F, Xu AJ (2010) Regulation of spinal neuroimmune responses by prolonged morphine treatment in a rat model of cancer induced bone pain. *Brain Res* 1326: 162-173.
- Cleeland CS, Bennett GJ, Dantzer R (2003) Are the symptoms of cancer and cancer treatment due to a shared biologic mechanism? *Cancer* 97: 2919-2925.
- Sang CN, Bennett GJ (2009) Novel therapies for the control and prevention of neuropathic pain. *Neuro Therapeutics* 6: 607-608.
- Dib-Hajj SD, Waxman SG (2014) Translational pain research: Lessons from genetics and genomics. *Sci Transl Med* 6: 249-244.
- Shenoy PA, Kuo A, Vetter I (2016) The walker 256 breast cancer cell-induced bone pain model in rats. *Front. Pharmacol* 7.
- Shenoy P, Kuo A, Vetter I (2017) Optimization and *in vivo* profiling of a refined rat model of walker 256 breast cancer cell-induced bone pain using behavioral, radiological, histological, immunohistochemical and pharmacological methods. *Front Pharmacol* p. 8.
- Shenoy PA, Kuo A, Khan N (2018) The somatostatin receptor-4 Agonist J-2156 alleviates mechanical hypersensitivity in a rat model of breast cancer induced bone pain. *Front Pharmacol* 9.
- Shenoy PA (2018) Establishment, optimization and characterization of a rat model of breast cancer induced bone pain. Faculty of Medicine, The University of Queensland, Australia.
- Kambiz S, Baas M, Duraku LS (2014) Innervation mapping of the hind paw of the rat using Evans Blue extravasation, Optical Surface Mapping and CASAM. *J Neurosci Methods* 229: 15-27.
- Cobianchi S, De Cruz J, Navarro X (2014) Assessment of sensory thresholds and nociceptive fiber growth after sciatic nerve injury reveals the differential contribution of collateral reinnervation and nerve regeneration to neuropathic pain. *Exp Neurol* 255: 1-11.
- De Koning P, Brakkee JH, Gispens WH (1986) Methods for producing a reproducible crush in the sciatic and tibial nerve of the rat and rapid and precise testing of return of sensory function. *J Neurol Sci* 74:237-246.
- Yang CJ, Wang XW, Li X (2011) A rat model of bone inflammation-induced pain by intra-tibial complete Freund's adjuvant injection. *Neurosci Lett* 490: 175-179.
- Tena B, Escobar B, Arguis MJ (2012) Reproducibility of electronic Von Frey and Von Frey monofilaments testing. *Clin J Pain* 28: 318-323.
- Reitz MC, Hrnčić D, Treede RD (2016) A comparative behavioural study of mechanical hypersensitivity in 2 pain models in rats and humans. *Pain* 157: 1248-1258.
- Abrams DI, Jay CA, Shade SB (2007) Cannabis in painful HIV-associated sensory neuropathy: A randomized placebo-controlled trial. *Neurol* 68: 515-521.
- Prabhavathi K, Chandra US, Soanker R (2014) A randomized, double blind, placebo controlled, cross over study to evaluate the analgesic activity of *Boswellia serrata* in healthy volunteers using mechanical pain model. *Indian J Pharmacol* 46: 475-479.
- Levine JD, Gordon NC, Taiwo YO (1988) Potentiation of pentazocine analgesia by low-dose naloxone. *J Clin Invest* 82: 1574-1577.
- Neuvonen PJ, Tokola O, Toivonen ML (1985) Methionine in paracetamol tablets, a tool to reduce paracetamol toxicity. *Int J Clin Pharmacol Ther* 23: 497-500.
- Staud R, Weyl EE, Price DD (2012) Mechanical and heat hyperalgesia highly predict clinical pain intensity in patients with chronic musculoskeletal pain syndromes. *J Pain* 13: 725-735.
- Scott AC, McConnell S, Laird B (2012) Quantitative sensory testing to assess the sensory characteristics of cancer-induced bone pain after radiotherapy and potential clinical biomarkers of response. *Eur J Pain* 16: 123-133.
- Fang D, Kong L-Y, Cai J (2015) Interleukin-6-mediated functional upregulation of TRPV1 receptors in dorsal root ganglion neurons through the activation of JAK/PI3K signaling pathway: Roles in the development of bone cancer pain in a rat model. *Pain* 156: 1124-1144.
- Liu M, Yang H, Fang D (2013) Upregulation of P2X3 receptors by neuronal calcium sensor protein VILIP-1 in dorsal root ganglions contributes to the bone cancer pain in rats. *Pain* 154: 1551-1568.
- Zheng Q, Fang D, Liu M (2013) Suppression of KCNQ/M (Kv7) potassium channels in dorsal root ganglion neurons contributes to the development of bone cancer pain in a rat model. *Pain* 154: 434-448.
- Shih MH, Kao SC, Wang W (2012) Spinal Cord NMDA receptor-mediated activation of mammalian target of rapamycin is required for the development and maintenance of bone cancer-induced pain hypersensitivities in rats. *J Pain* 13: 338-349.
- Li TF, Fan H, Wang YX (2016) Aconitum-derived bulleyaconitine A exhibits antihypersensitivity through direct stimulating dynorphin A expression in spinal microglia. *J Pain* 17: 530-548.
- Li Y, Cai J, Han Y (2014) Enhanced function of TRPV1 via up-regulation by insulin-like growth factor-1 in a rat model of bone cancer pain. *Eur J Pain* 18: 774-784.
- Liu S, Liu YP, Yue DM (2014) Protease-activated receptor 2 in dorsal root ganglion contributes to peripheral sensitization of bone cancer pain. *Eur J Pain* 18: 326-337.
- Zhang R-X, Li A, Liu B (2008) Electroacupuncture attenuates bone cancer-induced hyperalgesia and inhibits spinal preprodynorphin expression in a rat model. *Eur J Pain* 12: 870-878.
- Muralidharan A, Wyse BD, Smith MT (2014) Analgesic efficacy and mode of action of a selective small molecule angiotensin ii type 2 receptor antagonist in a rat model of prostate cancer-induced bone pain. *Pain Med* 15: 93-110.
- Wang LN, Yao M, Yang JP (2011) Cancer-induced bone pain sequentially activates the ERK/MAPK pathway in different cell types in the rat spinal cord. *Mol Pain* 7: 48.
- Pan HL, Zhang YQ, Zhao ZQ (2010) Involvement of lysophosphatidic acid in bone cancer pain by potentiation of TRPV1 via PKCε pathway in dorsal root ganglion neurons. *Mol Pain* 6: 85.
- Chen G, Kim YH, Li H (2017) PD-L1 inhibits acute and chronic pain by suppressing nociceptive neuron activity via PD-1. *Nat Neurosci* 20: 917-926.

- 36 De Felice M, Lambert D, Holen I (2016) Effects of Src-kinase inhibition in cancer-induced bone pain. *Mol Pain* 12: 174480691664372.
- 37 Cheng W, Chen YL, Wu L (2016) Inhibition of spinal UCHL1 attenuates pain facilitation in a cancer-induced bone pain model by inhibiting ubiquitin and glial activation. *Am J Transl Res* 8: 3041-3048.
- 38 Yao M, Chang XY, Chu YX (2011) Antiallodynic effects of propentofylline Elicited by interrupting spinal glial function in a rat model of bone cancer pain. *J Neurosci Res* 89: 1877-1886.
- 39 Sima L, Fan B, Yan L (2016) Effects of electroacupuncture treatment on bone cancer pain model with morphine tolerance. Evidence-based complementary and alternative medicine 1-5.
- 40 Pan HL, Liu BL, Lin W (2016) Modulation of Nav1.8 by lysophosphatidic acid in the induction of bone cancer pain. *Neurosci Bull* 32: 445-454.
- 41 Fu Q, Shi D, Zhou Y (2016) MHC-I promotes apoptosis of GABAergic interneurons in the spinal dorsal horn and contributes to cancer induced bone pain. *Exp Neurol* 286: 12-20.
- 42 Luo J, Huang X, Li Y (2016) GPR30 disrupts the balance of GABAergic and glutamatergic transmission in the spinal cord driving to the development of bone cancer pain. *Oncotarget* 7: 73462-73472.
- 43 Huang Q, Mao XF, Wu HY (2016) Bullatine A stimulates spinal microglial dynorphin A expression to produce anti-hypersensitivity in a variety of rat pain models. *J Neuroinflammation* 13: 214.
- 44 Du JY, Liang Y, Fang JF (2016) Effect of systemic injection of heterogenous and homogenous opioids on peripheral cellular immune response in rats with bone cancer pain: A comparative study. *Exp Ther Med* 12: 2568-2576.
- 45 Wang SF, Dong CG, Yang X (2016) Upregulation of (C-X-C motif) Ligand 13 (CXCL13) attenuates morphine analgesia in rats with cancer-induced bone pain. *Med Sci Monit* 22: 4612-4622.
- 46 Liu M, Yao M, Wang H (2017) P2Y12 receptor-mediated activation of spinal microglia and p38MAPK pathway contribute to cancer-induced bone pain. *J Pain Res* 10: 417-426.
- 47 Liu L, Gao XJ, Ren CG (2017) Monocyte chemoattractant protein-1 contributes to morphine tolerance in rats with cancer-induced bone pain. *Exp Ther Med* 13: 461-466.
- 48 Song Z, Xiong B, Zheng H (2017) STAT1 as a downstream mediator of ERK signaling contributes to bone cancer pain by regulating MHC II expression in spinal microglia. *Brain Behav Immun* 60: 161-173.
- 49 Yao W, Zhao H, Shi R (2017) Recombinant protein transduction domain-Cu/Zn superoxide dismutase alleviates bone cancer pain via peroxiredoxin 4 modulation and antioxidation. *Biochem Biophys Res Commun*. 486: 1143-1148.
- 50 Sun Y, Wu YX, Zhang P (2017) Anti-rheumatic drug iguratimod protects against cancer-induced bone pain and bone destruction in a rat model. *Oncol Lett* 13: 4849-4856.
- 51 Sun Y, Tian Y, Li H (2017) Antinociceptive effect of intrathecal injection of genetically engineered human bone marrow stem cells expressing the human proenkephalin gene in a rat model of bone cancer pain. *Pain Res Manag* 2017: 11.
- 52 Guo G, Peng Y, Xiong B (2017) Involvement of chemokine CXCL11 in the development of morphine tolerance in rats with cancer-induced bone pain. *J Neurochem* 141: 553-564.
- 53 Dai WL, Yan B, Jiang N (2017) Simultaneous inhibition of NMDA and mGlu1/5 receptors by levo-Corydalmine in rat spinal cord attenuates bone cancer pain. *Int J Cancer* 141: 805-815.
- 54 Hou X, Weng Y, Ouyang B (2017) HDAC inhibitor TSA ameliorates mechanical hypersensitivity and potentiates analgesic effect of morphine in a rat model of bone cancer pain by restoring mu-opioid receptor in spinal cord. *Brain Res* 1669: 97-105
- 55 Hang LH, Li SN, Dan X (2017) Involvement of Spinal CCR5/PKCgamma Signaling Pathway in the Maintenance of Cancer-Induced Bone Pain. *Neurochem Res* 42: 563-571.
- 56 Liang Y, Du JY, Fang JF (2017) Alleviating mechanical allodynia and modulating cellular immunity contribute to electro-acupuncture's dual effect on bone cancer pain. *Integr Cancer Ther* 17: 401-410.
- 57 Zhou YQ, Liu DQ, Chen SP (2018) Reactive oxygen species scavengers ameliorate mechanical allodynia in a rat model of cancer-induced bone pain. *Redox Biol* 14: 391-397.
- 58 Hu XF, He XT, Zhou KX (2017) The analgesic effects of triptolide in the bone cancer pain rats via inhibiting the upregulation of HDACs in spinal glial cells. *J Neuroinflammation* 14: 213.
- 59 Wang Y, Ni H, Li H (2018) Nuclear factor kappa B regulated monocyte chemoattractant protein-1/chemokine CC motif receptor-2 expressing in spinal cord contributes to the maintenance of cancer-induced bone pain in rats. *Mol Pain* 14: 1744806918788681.
- 60 Liang Y, Du JY, Fang JF (2018) Alleviating mechanical allodynia and modulating cellular immunity contribute to electroacupuncture's dual effect on bone cancer pain. *Integr Cancer Ther* 17: 401-410.
- 61 Guedon JMG, Longo G, Majuta LA (2016) Dissociation between the relief of skeletal pain behaviors and skin hypersensitivity in a model of bone cancer pain. *Pain* 157: 1239-1247.
- 62 Sabino MA, Luger NM, Mach DB (2003) Different tumors in bone each give rise to a distinct pattern of skeletal destruction, bone cancer-related pain behaviors and neurochemical changes in the central nervous system. *Int J Cancer* 104: 550-558.
- 63 Lozano Ondoua AN, Symons Liguori AM, Vanderah TW (2013) Cancer-induced bone pain: Mechanisms and models. *Neurosci Lett* 557: 52-59.
- 64 Caraceni A, Davies A, Poulain P (2013) Guidelines for the management of breakthrough pain in patients with cancer. *J Natl Compr Canc Netw* 1: S29-S36.
- 65 Laird BJ, Walley J, Murray GD (2011) Characterization of cancer-induced bone pain: an exploratory study. *Support Care Cancer* 19: 1393-1401.
- 66 Mogil JS, Graham AC, Ritchie J (2010) Hypolocomotion, asymmetrically directed behaviors (licking, lifting, flinching, and shaking) and dynamic weight bearing (gait) changes are not measures of neuropathic pain in mice. *Mol Pain* 6: 34.
- 67 Coulthard P, Simjee SU, Pleuvry BJ (2003) Gait analysis as a correlate of pain induced by carrageenan intraplantar injection. *J Neurosci Methods* 128: 95-102.
- 68 Piesla MJ, Leventhal L, Strassle BW (2009) Abnormal gait, due to inflammation but not nerve injury, reflects enhanced nociception in preclinical pain models. *Brain Res* 1295: 89-98.
- 69 Gabriel AF, Marcus MAE, Walenkamp GHIM (2009) The CatWalk method: Assessment of mechanical allodynia in experimental chronic pain. *Behav Brain Res* 198: 477-480.
- 70 Fernihough J, Gentry C, Malcangio M (2004) Pain related behaviour in two models of osteoarthritis in the rat knee. *Pain* 112: 83-93.
- 71 Boettger MK, Weber K, Schmidt M (2009) Gait abnormalities

- differentially indicate pain or structural joint damage in monoarticular antigen-induced arthritis. *Pain* 145: 142-150.
- 72 Aihara K, Ono T, Umei N (2017) A study of the relationships of changes in pain and gait after tourniquet-induced ischemia-reperfusion in rats. *J Phys Ther Sci* 29: 98-101.
- 73 Vlaeyen JWS, Linton SJ (2000) Fear-avoidance and its consequences in chronic musculoskeletal pain: A state of the art. *Pain* 85: 317-332.
- 74 Feinstein AB, Sturgeon JA, Darnall BD (2017) The effect of pain catastrophizing on outcomes: A developmental perspective across children, adolescents, and young adults with chronic pain. *J Pain* 18: 144-154.
- 75 Fingleton C, Smart K, Moloney N (2015) Pain sensitization in people with knee osteoarthritis: a systematic review and meta-analysis. *Osteoarthr Cartil* 23: 1043-1056.
- 76 Somers TJ, Keefe FJ, Pells JJ (2009) Pain catastrophizing and pain-related fear in osteoarthritis patients: relationships to pain and disability. *J Pain Symptom Manage* 37: 863-872.
- 77 Keefe FJ, Lefebvre JC, Egert JR (2000) The relationship of gender to pain, pain behavior, and disability in osteoarthritis patients: the role of catastrophizing. *Pain* 87: 325-334.
- 78 Tichonova A, Rimdeikienė I, Petruševičienė D (2016) The relationship between pain catastrophizing, kinesiophobia and subjective knee function during rehabilitation following anterior cruciate ligament reconstruction and meniscectomy: A pilot study. *Medicina* 52: 229-237.
- 79 Niederstrasser NG, Meulders A, Meulders M (2015) Pain catastrophizing and fear of pain predict the experience of pain in body parts not targeted by a delayed-onset muscle soreness procedure. *J Pain* 16: 1065-1076.
- 80 Haddas R, Lieberman IH, Block A (2018) The relationship between fear-avoidance and objective biomechanical measures of function in patients with adult degenerative scoliosis. *Spine* 43: 647-653
- 81 Al-Obaidi SM, Al-Zoabi B, Al-Shuwaie N (2003) The influence of pain and pain-related fear and disability beliefs on walking velocity in chronic low back pain. *Int J Rehabil Res* 26: 101-108.
- 82 Vincent HK, Adams MC, Vincent KR (2013) Musculoskeletal pain, fear avoidance behaviors, and functional decline in obesity: potential interventions to manage pain and maintain function. *Reg Anesth Pain Med* 38: 481-491.