

## A Case of Rectal Villous Adenoma in a 28-Year-Old Female Patient Presenting with Chronic Diarrhea and Acute Renal Failure

Sheref A Elseidy<sup>1\*</sup>, Mohammad A Khaskia<sup>1</sup>, Aya Madboly<sup>2</sup> and Haitham H Alzamli<sup>1</sup>

<sup>1</sup>Department of Tropical and Hepatology, Ain Shams University, Abbassia, Cairo, Egypt

<sup>2</sup>Department of Nutrition and Endocrinology, National Institute of Diabetes, Cairo, Egypt

**Corresponding author:** Sheref A. Elseidy, Senior Consultant, Department of Tropical and Hepatology, Faculty of Medicine, Ain Shams University, Abbassia 11566, Cairo, Egypt, Tel: +2001001571545; E-mail: drsheref.elseidy@gmail.com

**Rec Date:** November 24, 2017, **Acc Date:** November 28, 2017, **Pub Date:** November 30, 2017

**Citation:** Elseidy SA, Khaskia MA, Madboly A, Alzamli HH (2017) A Case of Rectal Villous Adenoma in A 28-Year-Old Female Patient Presenting with Chronic Diarrhea and Acute Renal Failure. J Emerg Intern Med Vol.1 No.2:15.

### Abstract

**Background:** Secretory villous adenoma is one of the rare causes of severe secretory diarrhea, acute kidney injury and refractory electrolyte imbalance is the classic triad of this disorder named McKittrick-Wheelock syndrome. Aggressive hydroelectrolyte rebalancing is often needed, and curative treatment is obtained only with complete removal of the lesion, by endoscopic resection or surgery. High clinical suspicion in patients with severe dehydration due secretory diarrhea and concomitant electrolyte imbalance is both mandatory and essential for early diagnosis and treatment before complications arise.

**Case presentation:** In this case we describe a 28-year-old female patient suffering from 6 months of secretory diarrhea, presented to our emergency department with severe dehydration, acute renal failure, hyponatremia and hypokalemia. After rehydration and electrolyte replacement confirmation, the diagnosis was done by colonoscopy and biopsy showing 3 sessile colonic polyps (2 at distal end of Transverse colon, largest one at descending colon) the biopsy confirmed the presence of villous adenoma with high grade dysplasia. Normalization of the electrolyte imbalance and the kidney functions was done, and she was successfully treated with surgical resection that lead to complete resolution of symptoms.

**Conclusion:** McKittrick-Wheelock syndrome is a rare life-threatening depletion syndrome that can lead to acute kidney injury and electrolyte imbalance with concomitant organ damage. Early management by vigorous repletion therapy and curative surgical resection will mostly resolve the symptoms and complications.

**Keywords:** Villous adenoma; Chronic diarrhea; Electrolyte imbalance; Acute renal failure tract

### Introduction

Secretory villous adenoma is one of the rare causes of severe secretory diarrhea, acute kidney injury and refractory electrolyte imbalance is the classic triad of this disorder named McKittrick-Wheelock syndrome. Aggressive hydroelectrolyte rebalancing is often needed, and curative treatment is obtained only with complete removal of the lesion, by endoscopic resection or surgery. High clinical suspicion in patients with severe dehydration due secretory diarrhea and concomitant electrolyte imbalance is both mandatory and essential for early diagnosis and treatment before complications arise [1-5].

In this case we describe a 28 year old female patient suffering from 6 months of secretory diarrhea, presented to our emergency department with severe dehydration, acute renal failure, hyponatremia and hypokalemia [6-9].

### Case Presentation

Our study reports a 28-year-old female patient presented to our ER with severe weakness, tachycardia and a 6-month history of non-bloody mucoid diarrhea with 5-6 motions/d, not improved with fasting, several courses of antibiotics.

- There was significant un-estimated weight loss; along with mild epigastric pain.
- Examination showed pallor, hypotension, HR: 120, bl.pr: 80/40.
- Labs showed BUN: 84, Creatinine: 1.1, HCT: 63, K: 1.8, Na: 109 and TLC: 9.1.
- ESR was found to be 30 and CRP < 6 with negative tumor markers with values: AFP: 3.1, CEA: 1.1, CA 19-9: 5.2 and CA 125: 9.7.
- Extensive volume repletion was done after admission with ringer and potassium infusion which lead to resolution of both clinical signs and symptoms, elevation of blood pressure and resolution of tachycardia with improvement of serum sodium and potassium with follow up results of (Na: 128 then 135, K: 3.1, 3.3, 4.3).
- PAUS: Liver of average size, homogenous and with smooth borders and 2 cm gall bladder stone.

- CT abdomen and pelvis showed hepatomegaly 17 cm (Homogenous CT density, no HFLs), Single GB stone (2 cm, normal GB wall thickness) and Multiple uterine fibroids.
- Colonoscopy was done and revealed 3 sessile colonic polyps (2 at distal end of Transverse colon, largest one at descending colon) with apparently normal mucosa in-between. Multiple biopsies were taken. Microscopic picture: Polypoid colonic mucosa showing villoglandular/glandular architecture with marked nuclear stratification of covering epithelium and the diagnosis of villous adenoma with high grade dysplasia was achieved.
- Volume and electrolyte repletion was continued till surgical intervention. After clinical and laboratory improvement the patient underwent endoscopic excision of the polyps which alleviated her symptoms.

## Discussion

Electrolyte depletion syndrome is a rare disorder caused by fluid and electrolyte hypersecretion from a villous adenoma. Dehydration with severe hyponatremia, hypokalemia, metabolic acidosis, acute renal failure and watery diarrhea is typical. Exact diagnosis is difficult require a high level of suspicion and early colonoscopy for visualization and biopsy. Most cases present at late stages mostly due to being misdiagnosed as infectious diarrhea as in this case. Villous adenomas rarely cause a secretory diarrhea syndrome and only about 3% of villous adenomas have secretory activity [1-9]. The tumor usually is located at the rectosigmoid or rectum and often is 3-4 cm or more in diameter [9,10], also secretory villous adenomas associated with depletion syndrome are large, ranging from 7 cm to 18 cm with increased diarrheal volumes of 350-3000 mL that leads to hypovolemia, metabolic imbalances and acute renal failure. The two main determinants of whether the adenoma would produce depletion syndrome are the large size which provides higher surface area for secretion, and the distal location which spares only a small area of normal mucosa remains for fluid absorption [11]. Complications are grave and can be prevented by adequate and prompt rehydration and acid base correction is not immediately performed and patient is stabilized [7,8].

Complications range from dehydration, asthenia and tachycardia up to profound acute renal injury and myocardial affection in the form of fatal arrhythmias [1,2]. it was found that up to 80 % of these villous adenomas have been reported to host foci of adenocarcinoma, making the most appropriate treatment is complete surgical excision of the lesion to decrease the chance of cancer and to resolve the depletion syndrome, also Surgical polypectomy is advised if the polyp is larger than 3 cm or sessile as villous adenomas often are [12].

In this case we discussed a young aged patient presented with this depletion syndrome confirmed to be owed to her rectal villous adenoma. to our knowledge this is the first case to present with this rare disorder at her age group. Management with early rehydration and electrolyte correction

stabilized the patient and it was followed by surgical resection that completely resolved her symptoms.

## Conclusion

A triad of acute prerenal failure, electrolyte disturbances and chronic diarrhea combined with the absence of other major causes of prolonged diarrhea should raise the indication for an early complete colonoscopy for prompt diagnosis in typical cases and to intervene with an appropriate therapy, resolving a condition that may develop life-threatening complications and that, if left untreated for a long time, could evolve in invasive cancer.

## References

1. McKittrick LS, Wheelock FC (1997) Carcinoma of the colon. *Dis Colon Rectum* 40: 1494-1495.
2. Choi WH, Ryuk J, Kim HJ, Park SY, Park JS, et al. (2012) A case of giant rectal villous tumor with severe fluid-electrolyte imbalance treated by laparoscopic low anterior resection. *J Korean Surg Soc* 82: 325-329.
3. McCabe RE, Kane KK, Zintel HA, Pierson RN (1970) Adenocarcinoma of the colon associated with severe hypokalemia: Report of a case. *Ann Surg* 172: 970-974.
4. Older J, Older P, Colker J, Brown R (1999) Secretory villous adenomas that cause depletion syndrome. *Arch Intern Med* 159: 879-880.
5. Targarona EM, Hernandez PM, Balague C, Martinez C, Hernández J, et al. (2008) McKittrick-Wheelock syndrome treated by laparoscopy: report of 3 cases. *Surg Laparosc Endosc Percutan Tech* 18: 536-538.
6. Konishi S, Nakada I, Satani T, Kasuga T, Watanabe Y, et al. (2010) Villous rectal adenoma as a rare cause of acute renal failure (McKittrick-Wheelock syndrome). *Coloproctology* 30: 251-253.
7. Popescu A, Orban-Schiopu AM, Becheanu G, Diculescu M (2005) McKittrick-Wheelock syndrome - A rare cause of acute renal failure. *Rom J Gastroenterol* 14: 63-66.
8. Emrich J, Niemeyer C (2002) The secreting villous adenoma as a rare cause of acute renal failure. *Med Klin (Munich)* 97: 619-623.
9. Watari J, Sakurai J, Morita T, Yamasaki T, Okugawa T, et al. (2011) A case of Cronkhite-Canada syndrome complicated by McKittrick-Wheelock syndrome associated with advanced villous adenocarcinoma. *Gastrointest Endosc* 73: 624-626.
10. Older J, Older P, Colker J, Brown R (1999) Secretory villous adenomas that cause depletion syndrome. *Arch Intern Med* 159: 879-880.
11. Lee YS, Lin HJ, Chen KT (2010) McKittrick-Wheelock syndrome: A rare cause of life-threatening electrolyte disturbances and volume depletion. *J Emerg Med*.
12. Agnes A, Novelli D, Doglietto GB, Valerio P (2016) A case report of a giant rectal adenoma causing secretory diarrhea and acute renal failure: McKittrick-Wheelock syndrome. *BMC Surgery* 16: 39.