

A young female presented with features of Gastric Outlet Obstruction

M.M.Shahin-Ul-Islam

Dhaka Medical College, Bangladesh

Abstract

Peutz-Jeghers syndrome (PJS) is an autosomal dominant syndrome characterized by multiple hamartomatous polyps in the gastrointestinal tract, mucocutaneous pigmentation and an increased risk of gastrointestinal cancer. Polyps are found throughout the gastrointestinal tract but commonly in the small bowel (60-90%), colon (50-64%), stomach (15-30%) and presents with gastrointestinal bleeding, anaemia, abdominal pain due to intussusceptions, and obstruction of small intestine or infarction. PJS as a cause of gastric outlet obstruction is rare. A young female of 13 years presented with recurrent attack of abdominal pain, vomiting with significant weight loss for last 4 years. Pain was severe, colicky, in epigastric, periumbilical region aggravated after taking food, relieved with vomiting (spontaneous and induced), vomituous contained 2-3 previous days undigested food. She lost 8 kg weight in last 1 year. She underwent appendectomy 3 years back. There was no history of fever, jaundice, haematemesis, melaena, alteration of bowel habit. She was moderately anaemic, nonicteric, BMI 15 kg/m² with black pigmentation in lower lip, mucous membrane, at finger tips and sole of foot (figure:1) with an ill-defined lump in epigastric region. Other systemic examination revealed normal. Her haemoglobin was 9 gm/dl, ESR, CRP, LFT, creatinine, CXR were normal with hyponatremia (Na 125), hypokalemia (K 2.5). USG of abdomen suggested gastric mass. After one unit of blood transfusion and correction of electrolyte imbalance, UGI endoscopy revealed multiple gastric & duodenal polyps with largest one was situated at lower body along the lesser curvature with a long pedicle, body of the polyp was impacted into pylorus (figure 2). Endoscopic polypectomy (figure 2) was done for larger polyps of stomach & duodenum, relieved obstruction with significant symptomatic improvement. Histopathology revealed hamartomatous polyp. Later on colonoscopy also revealed multiple polyps of variable sizes followed by colonoscopic polypectomy, histologically hamartomatous polyps. Our final diagnosis was PJS with gastric outlet obstruction due to gastric polyp with polyposis coli.

Received: January 07, 2022; **Accepted:** January 17, 2022; **Published:** January 28, 2022

Biography

M.M.Shahin-Ul-Islam, graduated from Dhaka Medical College, the best medical college in Bangladesh. He started his carrier in medicine and completed fellowship in internal medicine from Bangladesh College of Physicians and Surgeons (BCPS). He worked as consultant of internal medicine in district hospital of Bangladesh. After completing his specialization in Gastroenterology, he started working as assistant professor in Faridpur Medical College Hospital from

2010. Besides teaching undergraduate & post graduate students, he started Gastroenterology department at Faridpur Medical College in 2010. From last year working as associate professor (current charge) at same institute. He is a guide of a number of thesis and dissertations. He has a number of publications in the field of medicine and gastroenterology in national and international journals.