

## DIAGNOSTIC ACCURACY OF UTERINE ARTERY DOPPLER ULTRASOUND TO PREDICT PREECLAMPSIA IN PRIMIPAROUS FEMALES

**Mrs. Tahmina Yousaf**

Department of Medical Biochemistry, Govt. Medical college, Amritsar

**Objective:** To compare the outcome of oral misoprostol (20µg 2 hourly) with vaginal misoprostol (25µg 6 hourly) for induction of labour in term pregnancies, in term of frequency of vaginal delivery and cesarean section All pregnant women at term (>37 weeks gestation) with obstetric and medical indication for induction of labour and having Bishop Score  $\leq 6$  were included in this study. Parity  $\geq 4$ , previous history of obstetric and gynecological surgery and suspected cephalo pelvic disproportion were excluded. The study was conducted after approval from the ethical committee of the hospital. The subjects fulfilling inclusion criteria were enrolled after informed consent. hours interval up to a maximum of 4 doses. Randomization was done using lottery method. Data was entered on predesigned performa and was analyzed using SPSS version



Tahmina Yousaf . MBBS(Pb), MCPS.FCPS(Pk)  
Associate Gynecologist.

Federal General Hospital , Chak Shahzad, Islamabad Dr.Tahmina Yousaf is a consultant Obstetrician and Gynaecologist practicing at Medicsi, Islamabad and Care plus, Islamabad. She practices between Monday to Thursday. Now you can easily get an online appointment with Dr Tahmina Yousaf through Marham

### Publications:

Anil Batta, Nitric Oxide mechanism in the protective effect of naringin against post stroke depression

**Key words:** Induction of Labour, misoprostol, oral and vaginal route.

[4<sup>th</sup> international conference on Ophthalmology, August 07-08, 2020, Osaka, Japan](#)

**Abstract Citation:** [Anil Batta, Scientific production in Neuroophthalmology, World ophthalmology 2020, august 07-08, 2020, Osaka, Japan](#)

