

Efficiency and economics of weed control with pre and post-emergence herbicides in soybean (*Glycine Max* L.)

¹Sylvestre Habimana*, ¹K. N. Kalyana Murthy, ¹B. C. Shankaralingappa, ²M. T. Sanjay and ³C. Ramachandra

¹Department of Agronomy, University of Agricultural Sciences, GKVK- Bengaluru India

²All India Coordinated Research Project on Weed Control, Hebbal-Bengaluru India

³All India Coordinated Research Project on Rice, V.C Farm Mandya India

ABSTRACT

A field experiment was conducted in zonal agricultural research station, university of agricultural sciences, (UAS) GKVK Bengaluru during Kharif, 2012 to study efficacy of pre and post-emergence herbicides on yield and economics of soybean. The experiment was laid out in Randomized Complete Block Design (RCBD) with ten treatments, replicated thrice. At harvest, intercultivation fb hand weeding at 20 and 40 DAS was found effective for controlling grasses, broad leaved weeds and sedge weeds, registered higher grain (2570 kg ha⁻¹) and haulm yield (2964 kg ha⁻¹). However, intercultivation fb. hand weeding at 20 and 40 DAS remained at par with pre-emergence application of metribuzin 70 WP fb. imazethapyr as well as metribuzin 70 WP fb. intercultivation at 30 DAS as compared to unweeded check. Nevertheless, benefit cost ratio was recorded higher (3.55) under the metribuzin 70 WP fb. imazethapyr.

Keywords: Economics, pre and post-emergence herbicides, soybean, weeds, yield.

INTRODUCTION

Soybean (*Glycine max*), is an important oil-yielding rainy-season (*kharif*) crop having multiple uses. It has revolutionized the rural economy and has improved socio-economic status of the farmers. Soybean produces 2-3 times more high quality protein yield per hectare than other pulses and cholesterol free oil [1]. It is preferred especially by vegetarians on account of its richness in protein, fat, carbohydrates, mineral, salts and vitamins. Among the various factors responsible for the low yield of soybean, weeds have been considered to be of prime importance.

The losses caused by weeds exceed the losses from any other category of biotic factors like insects, nematodes, diseases, rodents, mites, etc. Soybean is mostly grown in *kharif* season and suffers from severe weed crop competition due to continuous rain, which do not permit hand weeding operation timely resulted in yield loss to the tune of 30–80 per cent [2].

In general, judicious use of herbicides in crop land generally increase crop yield, improve crop quality and reduce production costs [3]. Therefore, an experiment was planned to study the efficacy of pre and post emergence herbicides on yield and economics of *kharif* soybean.

MATERIALS AND METHODS

Field experiment was conducted in zonal agricultural research station, university of agricultural sciences, GKVK Bengaluru during Kharif, 2012 to study the efficacy of pre and post-emergence herbicides on yield and economics of *kharif* soybean. The soil of the experimental site was red sandy loam, with slightly acidic (pH 6.44), medium in organic carbon (0.55 %), medium in available Nitrogen (288.549 kg ha⁻¹), in available potassium (175.08 kg ha⁻¹)

and in phosphorus (38.49 kg ha^{-1}). The experiment was laid out in RCBD and composed of ten treatments replicated thrice *viz* T₁: Pendimethalin 30 EC at $0.75 \text{ kg a.i ha}^{-1}$ at 3 DAS (days after sowing), T₂: Pendimethalin 30 EC at $0.75 \text{ kg a.i ha}^{-1}$ at 3 DAS followed by (fb) intercultivation (IC) at 30 DAS, T₃: Pendimethalin 30 EC at $0.75 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. fenoxaprop-p-ethyl 9 EC at 70 g a.i ha^{-1} at 20 DAS, T₄: Pendimethalin 30 EC at $0.75 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. imazethapyr 10 SC at $100 \text{ g a.i ha}^{-1}$ at 20 DAS, T₅: Metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS, T₆: Metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. IC at 30 DAS, T₇: Metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. fenoxaprop-p-ethyl 9 EC at 70 g a.i ha^{-1} at 20 DAS, T₈: Metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. imazethapyr 10 SC at $100 \text{ g a.i ha}^{-1}$ at 20 DAS, T₉: Intercultivation fb. hand weeding (HW) at 20 and 40 DAS, T₁₀: Unweeded check. Variety MAUS-2 was sown on 12th August 2012. Recommended fertilizer dose of 25 kg N , $60 \text{ kg P}_2\text{O}_5$ and $25 \text{ kg K}_2\text{O ha}^{-1}$ was applied in the form of urea, single super phosphate and muriate of potash, respectively. The observations on weeds density and dry weight, yield and yield attributes were recorded at harvest. The economics of weed control was worked out based on agricultural inputs and outputs. The data were analyzed statistically for test of significance following the procedure described by [4]. The level of significance on "F" test was tested at 5 per cent. The interpretation of data was done by using CD values calculated at $p \geq 0.05$.

The major weed flora observed in the experimental plot was the following: *Eleusine indica*, *Digitaria marginata*, *Dactyloctenium aegyptium*, *Eragrostis pilosa*, *Amaranthus viridis*, *Oldenlandia corymbosa*, *Parthenium hysterophorus*, *Commelina benghalensis*, *Acanthospermum hispidum*, *Borreria hispida* and *Cyperus rotundus*.

RESULTS AND DISCUSSION

Effect of pre and post-emergence herbicides on weed growth:

At harvest, among the weed control treatments significantly lower total weed density was recorded under intercultivation fb two hand weedings at 20 and 40 DAS (8.00) and remained at par with metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. imazethapyr 10 SC at $100 \text{ g a.i ha}^{-1}$ at 20 DAS (8.67) (Table 1). Similarly, significantly lower total weed dry weight as well as more than 96 per cent weed control efficiency were recorded in intercultivation fb two hand weedings at 20 and 40 DAS ($1.00 \text{ g } 0.25 \text{ m}^2$). However, intercultivation fb two hand weedings at 20 and 40 DAS remained at par with metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. imazethapyr 10 SC at $100 \text{ g a.i ha}^{-1}$ at 20 DAS ($1.17 \text{ g } 0.25 \text{ m}^2$). Similar trend was also noticed by [5,6]. Unweeded check treatment registered significantly higher total weed population (125.33), dry weight of weeds ($30.72 \text{ g } 0.25 \text{ m}^2$) as well as lower weed control efficiency (0.00 %). Better weed control efficiency could be attributed to better efficacy of this pre and post-emergence herbicides used in controlling grasses, broad leaved weeds as well as sedge weeds, which is evidenced by lower number of weeds per 0.25 m^2 and weed dry weight at harvest.

Effect of pre and post-emergence herbicides on yield and economics:

Intercultivation fb two hand weedings at 20 and 40 DAS registered higher grain yield (2570 kg ha^{-1}), haulm yield (2964 kg ha^{-1}), and higher net returns (49652 Rs.ha^{-1}) which remained at par with metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. imazethapyr 10 SC at $100 \text{ g a.i ha}^{-1}$ at 20 DAS (2497 kg ha^{-1} , 2867 kg ha^{-1} and 49013 Rs.ha^{-1} , respectively) (Table 2). Unweeded check recorded lower grain yield (496 kg ha^{-1}) haulm yield (864 kg ha^{-1}) and lower negative returns (-2374 Rs.ha^{-1}).

Table 1: Effect of weed management practices on weed density, weed dry weight and weed control efficiency (WCE)

Weed management practices	Weed density (no./0.25m ²)	Weed dry weight (g 0.25m ²)	WCE (%)
T ₁ : Pendimethalin 30 EC at $0.75 \text{ kg a.i ha}^{-1}$ at 3 DAS (days after sowing)	5.29(27.67)	3.27(10.28)	66.55
T ₂ : Pendimethalin 30 EC at $0.75 \text{ kg a.i ha}^{-1}$ at 3 DAS followed by (fb) intercultivation (IC) at 30 DAS	3.26(10.67)	1.63(2.24)	92.72
T ₃ : Pendimethalin 30 EC at $0.75 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. fenoxaprop-p-ethyl 9 EC at 70 g a.i ha^{-1} at 20 DAS	4.81(22.67)	2.77(7.26)	76.36
T ₄ : Pendimethalin 30 EC at $0.75 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. imazethapyr 10 SC at $100 \text{ g a.i ha}^{-1}$ at 20 DAS	3.89(14.67)	2.45(5.54)	81.96
T ₅ : Metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS	4.91(23.67)	3.15(9.50)	69.08
T ₆ : Metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. IC at 30 DAS	3.18(9.67)	1.32(1.26)	95.91
T ₇ : Metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. fenoxaprop-p-ethyl 9 EC at 70 g a.i ha^{-1} at 20 DAS	4.34(18.33)	2.63(6.44)	79.05
T ₈ : Metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. imazethapyr 10 SC at $100 \text{ g a.i ha}^{-1}$ at 20 DAS	2.85(8.67)	1.29(1.17)	96.19
T ₉ : IC fb. HW at 20 and 40 DAS	2.90(8.00)	1.22(1.00)	96.75
T ₁₀ : Unweeded check	11.20(125.33)	5.55(30.72)	0.00
CD (P = 0.05)	0.88	0.60	-

Data within parentheses are original values; data analyzed using transformation $-\sqrt{x} + 0.5$

The crop yield is inversely related to the weed index, lower weed index was observed with intercultivation fb. hand weeding at 20 and 40 DAS (0.00) and among the herbicides treatments, metribuzin 70 WP at $0.5 \text{ kg a.i ha}^{-1}$ at 3 DAS fb. imazethapyr 10 SC at $100 \text{ g a.i ha}^{-1}$ at 20 DAS recorded lower weed index (2.84) as it could be attributed to lower dry weight of weeds per ha as well as weed density. Similar results were also reported by [7].

Moreover, benefit cost ratio was recorded higher (3.55) under the metribuzin 70 WP at 0.5 kg a.i ha⁻¹ at 3 DAS fb. imazethapyr 10 SC at 100 g a.i. ha⁻¹ at 20 DAS, followed by Metribuzin 70 WP at 0.5 kg a.i ha⁻¹ at 3 DAS fb. IC at 30 DAS (3.33) and intercultivation fb. hand weeding at 20 and 40 DAS (3.29) as compared to unweeded check (-0.87) (Table 2). This was mainly due to lesser cost of cultivation. These results are in conformity with [8].

Table 2: Effect of weed management practices on grain yield, haulm yield, net returns, benefit cost ratio and weed index of soybean

Weed management practices	Grain yield (kg ha ⁻¹)	Haulm yield (kg ha ⁻¹)	Net returns (Rs/ha)	Benefit cost ratio	Weed Index (%)
T ₁ : Pendimethalin 30 EC at 0.75 kg a.i ha ⁻¹ at 3 DAS (days after sowing)	1500	2422	26517	2.43	41.63
T ₂ : Pendimethalin 30 EC at 0.75 kg a.i ha ⁻¹ at 3 DAS followed by (fb) intercultivation (IC) at 30 DAS	2105	2669	40796	3.16	18.09
T ₃ : Pendimethalin 30 EC at 0.75 kg a.i ha ⁻¹ at 3 DAS fb. fenoxaprop-p-ethyl 9 EC at 70 g a.i ha ⁻¹ at 20 DAS	1767	2607	32180	2.63	31.24
T ₄ : Pendimethalin 30 EC at 0.75 kg a.i ha ⁻¹ at 3 DAS fb. imazethapyr 10 SC at 100 g a.i. ha ⁻¹ at 20 DAS	2087	2469	37878	2.86	18.79
T ₅ : Metribuzin 70 WP at 0.5 kg a.i ha ⁻¹ at 3 DAS	1733	2458	31408	2.65	32.56
T ₆ : Metribuzin 70 WP at 0.5 kg a.i ha ⁻¹ at 3 DAS fb. IC at 30 DAS	2491	2693	48501	3.33	3.07
T ₇ : Metribuzin 70 WP at 0.5 kg a.i ha ⁻¹ at 3 DAS fb. fenoxaprop-p-ethyl 9 EC at 70 g a.i ha ⁻¹ at 20 DAS	1850	2110	31109	2.55	28.01
T ₈ : Metribuzin 70 WP at 0.5 kg a.i ha ⁻¹ at 3 DAS fb. imazethapyr 10 SC at 100 g a.i. ha ⁻¹ at 20 DAS	2497	2867	49013	3.55	2.84
T ₉ : IC fb. HW at 20 and 40 DAS	2570	2964	49652	3.29	0.00
T ₁₀ : Unweeded check	496	864	-2374	-0.87	80.70
CD (P = 0.05)	405.72	421.12	-	-	-

CONCLUSION

In light of the results obtained from the present investigation, it was concluded that for effective weeds control and securing maximum grain yield of soybean as well as economic returns, intercultivation fb two hand weedings at 20 and 40 DAS or application of pre-emergence herbicide metribuzin 70 WP at 0.5 kg a.i ha⁻¹ at 3 DAS followed by a post-emergence herbicide imazethapyr 10 SC at 100 g a.i. ha⁻¹ at 20 DAS may be adopted. This herbicide weed management method found to be promising to control weeds in soybean crop and would play an important role in areas where labor is too expensive and time is a constraint.

Acknowledgement

I would like to thank the Indian Council for Agricultural Research (ICAR), Indo-African fellowship program and Department of Agronomy, UAS GKVK Bengaluru for providing all possible research facilities while executing the field experiment and laboratory analysis.

REFERENCES

- [1] Kumari K.V.S.M., Rajeswari B. and Reddy B.M. **2002**. *Indian J. Plant Prot.*, **30** (2): 139-143.
- [2] Yaduraju N.T. **2002**. Weed management in oilseed crops. In: *Oilseeds and Oils : Research and Development Needs*. (Eds.). *Indian Society of Oilseed Res.*, 172-183.
- [3] Balyan R.S. and Malik R.K. **2003**. *Indian J. Weed Sci.*, **35** (1&2): 62-66.
- [4] Gomez, K. A. and Gomez, A. A., **1984**. Statistical Procedures Agricultural Research, (2/E) An International Rice Research Institute Book, A Wiley Inter Science publication, *John Wiley and Sons, New York*.
- [5] Shete B. T., Deore, N. R., and Tambe, A. D., **2008**. *J. Maharashtra Agric. Univ.*, **33** (2): 266-267.
- [6] Tiwari, D. K., Kewat, M. L., Khan, J. A. and Khamparia, N. K., **2007**. *Indian J. Agron.*, **52** (1): 74-76.
- [7] Suresh Kumar., Angiras, N.N., Rana, S.S. and Arvind S.T., **2008**, *Indian J. Weed Sci.*, **40** (1&2): 34-38.
- [8] Vijayalaxmi G. S., S. M. Hiremath, J. A. Hosmath, P. L. Patil, **2012**. *Karnataka J. Agric. Sci.*, **25** (2): (262-263)