

Genetic Analysis of *Cannabis* Samples Using STR Markers

Emma Hu*

Department of Neuropathology, Tokyo Metropolitan Neurological Hospital, Fuchu, Tokyo, Japan

Corresponding author: Emma Hu, Department of Neuropathology, Tokyo Metropolitan Neurological Hospital, Fuchu, Tokyo, Japan, E-mail: emma@gmail.com

Received date: July 18, 2023, Manuscript No. IPGJRR-23-17753; **Editor assigned date:** July 21, 2023, PreQC No. IPGJRR-23-17753 (PQ); **Reviewed date:** August 04, 2023, QC No. IPGJRR-23-17753; **Revised date:** August 11, 2023, Manuscript No. IPGJRR-23-17753 (R); **Published date:** August 18, 2023, DOI: 10.36648/2393-8854.10.4.62

Citation: Hu E (2023) Genetic Analysis of *Cannabis* Samples Using STR Markers. Glob J Res Rev Vol .10 No.4:62.

Description

The American Public Health Association (APHA) standard protocols were followed for the measurement of biological, environmental, and physiological parameters. When the greatest number of COVID-19 cases was clinically diagnosed, SARS-CoV-2 viral shedding in wastewater reached its peak. Were found during the study, with WS representing and CS representing of the positive samples. *Cannabis* plants have been grown for thousands of years all over the world, and they are used a lot in the textile industry and medicine. They can be divided into two categories for analyses of the needs of the justice system: fibrous hemp and narcotic hemp, which contain different amounts of the psychoactive compound THC. The guidelines stipulate that narcotic varieties are those with a content of active compounds greater than to distinguish between these two kinds of *Cannabis*, toxicological analysis is the most common method. Genetic identification, on the other hand, might be a good option if there are issues with the type of plant part sent to the laboratory or the amount of differentiating active compounds present. The genetic analysis of *cannabis* samples using STR markers was the subject of the research consortium's investigation. The fundamental statistical parameters that are used to describe the population were taken into account in the analysis of the collected population database of *Cannabis* samples from fibrous and narcotic varieties.

Genetic Markers

On the basis of a selected set of 23 STR genetic markers, which were subjected to preliminary statistical analyses, it was used to develop a method for genetic profiling *Cannabis*. His highest CT value occurred when the number of COVID-19 patients was lowest. The genetic models that have been developed will be used to select the genetic markers that are most useful for the genetic identification of *Cannabis*, which will make it possible to develop a tool for forensic experts to identify the plant. Wastewater Based Epidemiology (WBE) is an important tool in the fight against COVID-19 because it provides cost-effective, rapid, and non-invasive insights into the health status of the targeted population—from a single house to a large municipality. The implementation of Wastewater-Based Surveillance (WBS) has the potential to alleviate pressure on the public health system, assist in the management of pandemics,

aid in the formulation of well-informed choices, and safeguard public health. This study aimed to monitor the prevalence of SARS-CoV-2 genetic markers in Clinical Specimens (CS) and Wastewater Samples (WS) for 30 days in a house with COVID-19 patients. SARS-CoV-2's non-structural and structural-Nucleocapsid (N) protein genes were targeted by RT-qPCR in accordance with a tested experimental protocol. WS was used to determine which copy number was the lowest. On the other hand, when the number of COVID-19 patients was highest, WS obtained the lowest CT value, or highest copy numbers. The COVID-19 patient's advance signal of an increased SARS-CoV-2 viral load was detected in the WS earlier than in the CS. Involving redid preliminary sets in a conventional PCR approach; we affirmed that all SARS-CoV-2 variations recognized in both CS and WS were Delta variations. This is the first follow-up study to our knowledge to determine a temporal relationship between COVID-19 patients' discharge of SARS-CoV-2 RNA genetic markers in wastewater from a single house that included all family members for clinical sampling from a developing nation (Bangladesh) that does not have a proper sewage system. The study's most important findings indicate that COVID-19 cases could be identified by monitoring the genetic markers of the SARS-CoV-2 virus in wastewater, easing the burden on the public health system during COVID-19 pandemics.

Genetic Structure

Using microsatellite markers, the current study aimed to reveal the genetic structure of this candidate species from the Bay of Bengal and the Arabian Sea. Twelve microsatellite markers utilized for hereditary stock portrayal and 384 examples gathered from six unique areas were utilized for the review. A single unit stock of red snapper was found in the sampling region, and microsatellite markers suggested that there was sufficient gene flow between populations to limit genetic structuring. An examination of the genetic differentiation coefficient suggested that there was no structuring between populations. However, structure's Bayesian cluster analysis revealed some evidence of sub-structuring between populations in the Arabian Sea and the Bay of Bengal. The aforementioned findings were supported in part by isolation by distance analysis, Principal Coordinate Analysis (PCoA), and hierarchical AMOVA analysis. Before appropriate management strategies for the mangrove red snapper in Indian waters can be developed, it is

necessary to use more potent markers like SNPs to reveal the sub-structuring because microsatellites were unable to detect population differentiation in this instance. Farmers' traditional rice varieties are useful genetic resources for future rice improvement. These cultivars are extremely well-suited to a wide range of agro-ecological conditions. The accessions had a lot of different genes. With an average of 6.2 alleles per locus, the total number of alleles ranged from 2 to 22. The mean effective number of alleles per locus was 1.86, ranging from the median estimate of gene diversity was 0.355, with estimates ranging from. The average Polymorphism Information Content (PIC) value was 0.36, with values ranging from various leveled bunching investigation assembled the genotypes into two significant bunches with two sub-gatherings, which related with

the geographic beginnings. The dwarf accessions were in the first cluster, and the tall accessions were all in the second cluster. With two major populations, the results of the structure analysis were in line with those of the cluster analysis. The findings suggested that marker-assisted selection and genetic diversity in coconut could be assessed using SSR markers. In the Indo-Pacific region, the mangrove red snapper, *Lutjanus argentimaculatus*, is a marine fish of significant economic and cultural significance. It is now regarded as a much more significant species for aquaculture than for capture fisheries. An assortment of 96 palms, addressing 16 different coconut promotions from the Worldwide Coconut Gene bank for South Asia and the center East, Kidu, India, was examined for hereditary variety and populace structure in view of 35 basic grouping rehash loci.