



## New angle of plant diseases and its hero is Melanospora chionea causing new disease whose called pink root rot in several hosts.

**F.M. Farag**

Plant Pathol Res. Inst., ARC, Egypt.

**Abstract:**

Symptoms of pink root rot were observed on tomato (*Lycopersicon esculentum* Mill.) grown in Beni Sweif governorate (Nasser, Sumosta, Beba and El-Wasta county) in summer 2013. Based on morphological characteristics of the isolated fungus, disease symptoms and a pathogenicity test, *M. chionea* was identified as the causal agent of pink root rot of tomato. *M. chionea* has not previously been reported on tomato. This disease has infected numerous hosts belong to Asteraceae, Brasicaceae, Cucurbitaceae, Fabaceae, Molvaceae and Solanaceae.

**Keywords:** Melanospora chionea, pink root rot, Morphological characteristic, pathogenicity test,

**Publication of speakers:**

1. Cannon, P. F. 1982. A re-evaluation of *Melanospora corda* and similar Pyrenomycetes, with a revision of the British species Botanical journal of the Linnean Sociep. 84: 115-160.
2. Cannon P.F., Buddie AG, Bridge PD. 2008. The typification of *Colletotrichum gloeosporioides*. Mycotaxon 104:189-204.
3. Chaudhary P., Campbell J, Hawksworth DL, Sastry KN. 2006. *Vittatispora*, a new melanoporous genus from Indian soil. Mycologia 98:460-467, doi:10.3852/mycolgia.98.3.460
4. Davey ML, Tsuneda A, Currah RS. 2008. Evidence that the gemmae of *Papulaspora sepedonioides* are neotenous perithecia in the Melanoporales. Mycologia 100:626-635, doi:10.3852/08-001R
5. Gams W, Diederich P, Po~ldmaa K. 2004. Fungicolous fungi. In: Mueller GM, Bills GF, Foster MS, eds. Biodiversity of Fungi: inventory and monitoring methods. Amsterdam, the Netherlands: Elsevier. p 343-392.
6. Garcí'a D, Stchigel AM, Guarro J. 2002. A new species of *Sympastospora* from tropical soils. Mycologia 94:862-865,
7. Garcí'a D, Stchigel AM, Guarro J 2004. Two new species of Sphaerodes from Spanish soils. Stud Mycol 50:63-68.
8. Hanllin, R.T., T.K.Goh, and O. Tortotera, 1993. Icones Aascomycetum Venezuelae *Melanospora zamiae* Mycotaxon 49:167-174.
9. Hawksworth DL, Henrici A, Kirk PM. 1999. *Melanospora pegleri*, a new species from fallen *Laurus* and other leaves in the British Isles. Kew Bull 54:795-799,
10. Jeffries, P. & Young, T. W. K. 1994. Interfungal Parasitic Relationships.CAB International, Wallingford
11. Krug JC. 1988. A new species of *Persicospora* from African soil. Mycologia 80:414-417, doi:10.2307/3807643.
12. Luttrell,E.S. (1951). Taxonomy of the pyrenomycetes. Univ. Missouri Stud. 24: 1-120.
13. Nitzan N, Rogers JD, Johnson DA. 2004. *Melanospora washingtonensis*, a new species from potato. Sydowia 56:281-287.
14. Stchigel AM, Guarro J, Figueras MJ. (1996). A new species of *Melanospora* from India. Mycologia 101:446-448.
15. Stchigel AM, Cano J, Guarro J. 1999. A new species of *Melanospora* from Easter Island. Mycol Res 103:1305- 1308, doi:10.1017/S095375629900845X
16. Vujanovic V, Goh YK. 2009. *Sphaerodes mycoparasitica* sp. nov., a new biotrophic mycoparasite on *Fusarium avanaceum*, *F. graminearum* and *F. oxysporum*. Mycol Res 113:1172-1180, doi:10.1016/j.mycres.2009.07.018

**Emerging Trends in Plant Science and Natural Products Research, March 19-20, 2020; London, UK**

**Citation:** Eftekhar Baranian Kabir; The effects of land-use/cover change on some soil physical and chemical properties in central Iran; Natural Products 2020; March 19-20, 2020; London, UK