

## Pediatric Intensive Care Unit Admissions from the Emergency Departments in India - The 2018 Academic College of Emergency Experts Consensus Recommendations

**Lalitha AV<sup>1\*</sup>, Bernhard Fassl<sup>2\*</sup>, Ramon E. Gist<sup>3</sup>, Binita R. Shah<sup>4</sup>, Nitin Chawla<sup>5</sup>, Ajay Singh<sup>6</sup>, Arun Baranawali<sup>7</sup>, Shivakumar Shamarao<sup>8</sup>, Raghavendra Vanaki<sup>9</sup>, Prashant Maharjan<sup>10</sup>, Vivek Chauhan<sup>11</sup>, Prerna Batra<sup>12</sup>, Abhijeet Saha<sup>13</sup>, Reena Patel<sup>14</sup>, Sagar Galwankar<sup>15</sup> and Santosh Soans<sup>16</sup>**

<sup>1</sup>Pediatric Intensive Care and Pediatric Emergency, St. John's Medical College and Hospital, Bangalore, India

<sup>2</sup>Department of Pediatrics, University of Utah, Salt Lake City, Utah, USA

<sup>3</sup>Pediatric Critical Care Medicine, Assistant Professor of Pediatric, SUNY Downstate Medical Center, Brooklyn, New York

<sup>4</sup>Distinguished Professor of Emergency Medicine and Pediatrics; SUNY Downstate Medical Center, Brooklyn, NY

<sup>5</sup>Acute and critical care consultant, Kamineni Super Speciality Hospital, Hyderabad, India

<sup>6</sup>Department of Pediatric Orthopedic and Trauma, King George's Medical University, Lucknow, UP, India

<sup>7</sup>Department of Pediatric Emergency and Intensive Care, PGIMER, Chandigarh, India

<sup>8</sup>Consultant Pediatric Intensivist, Manipal hospital, Bangalore, India

<sup>9</sup>Department of Pediatrics, SN Medical College, Bagalkot, India

<sup>10</sup>Department of Emergency Medicine, University of Michigan, Ann Arbor, United States

<sup>11</sup>Kangra University and Medical Center, Dharamsala, India

<sup>12</sup>Department of Pediatrics, University College of Medical Sciences and Guru Teg Bahadur Hospital, Delhi, India

<sup>13</sup>Department of Pediatrics, Lady Hardinge Medical College and Associated Kalawati Saran Children Hospital, New Delhi, India

<sup>14</sup>Department of Pediatrics, University of Utah, Salt Lake City, Utah, USA

<sup>15</sup>Emergency Medicine, University of Florida, Jacksonville, Florida

<sup>16</sup>Department of Pediatrics at AJ Institute of Medical Science and Neonatal and Pediatric Intensive Care Division, Mangalore, India

**\*Corresponding author:** Lalitha AV, Pediatric Intensive Care and Pediatric Emergency, St John's Medical College Hospital, Bangalore, India, Tel: +919448461673; E-mail: drlalitha03@gmail.com

Bernhard Fassl, Department of Pediatrics, University of Utah, Salt Lake City, Utah, USA, Tel: +1 (801) 662-3645; E-mail: Bernhard.fassl@hsc.utah.edu

**Received:** January 02, 2019; **Accepted:** January 23, 2019; **Published:** January 30, 2019

**Citation:** Lalitha AV, Fassl B, Gist R, Shah B, Chawla N, et al. (2019) Pediatric Intensive Care Unit Admissions from the Emergency Departments in India - The 2018 Academic College of Emergency Experts Consensus Recommendations. J Emerg Trauma Care Vol.4 No.1:2.

### Abstract

There are global variations in policies that define clear indications for PICU (Pediatric Intensive care unit) admissions. In resource limited countries where PICU service availability is limited, the admission criteria to PICU are urgently needed to optimize the utilization of available intensive care services and to maximize patient benefit. The objective of these consensus recommendations on PICU admission criteria is to provide a framework and reference for future policy development by professional societies and governments.

**Design:** The consensus recommendations were developed by a multidisciplinary consensus task force comprised of

international experts in pediatric critical care, emergency medicine, trauma, critical care, and health policy stakeholders during the 2016 annual INDUSEM WORLD CONGRESS in Bengaluru, India.

**Measurements and main results:** A task force steering committee completed a global literature search about PICU admission criteria development; reviewed PICU admission guidelines published by a variety of professional organizations worldwide, and performed a literature review of relevant publications. The objectives of this Task Force is to provide a framework for validated approach to determine appropriateness of ICU admission in India(resource limited setting) based on a) prioritization modeling; b) general clinical criteria; c) clinical and

objective parameters and d) other criteria. The expert consensus panel then discussed and ranked proposed criteria according to scientific evidence, current standard of care, and expert opinion in the context of the Indian health system. The general subject was addressed in sections: admission criteria and benefits of different levels of care, following the appraisal of the literature, discussion, and consensus, recommendations were written.

**Conclusion:** Although these are consensus recommendations, the subjects addressed encompass complex ethical and medico-legal aspects of patient care that affect daily clinical practice. The scarcity of high-quality evidence made it difficult to answer all the questions asked related to ICU admission. Despite these limitations, the members of the Task Force believe that these recommendations provide a comprehensive framework to guide practitioners in making informed decisions during the admission process. This publication is designed to assist in future development of health policies to ensure effective resource allocation, maximize healthcare benefits and improve access to quality care for children.

**Keywords:** Pediatric intensive care; Admission criteria; PICU; Consensus recommendations

## Introduction

The PICU concept was initially developed about 40 years ago with the first consensus conference on critical care admission held in 1983 by the National Institute of Health in the US [1,2]. The principle that emerged from this group continues to be relevant even today as it identifies patients who should be admitted to the PICU as those who “reversible medical conditions with a reasonable prospect of substantial recovery” [3,4]. As with any treatment, the decision to admit a patient to the PICU should be based on potential benefit [5]. Pediatric intensive care admission criteria should select those patients who are the most likely to benefit from this level of care. Such patients are generally those who are severely ill and unstable, with a high likelihood of functional recovery after treatment of the acute illness [6,7]. Identification of patients who are “too well” or “too severely ill” for PICU admission is a complicated task and may be difficult if decisions are solely based upon diagnosis. Similarly, severity of illness scores such as the Pediatric Risk of Mortality Score (PRISM), Acute Physiology and Chronic Health Evaluation (APACHE), and Simplified Acute Physiology Scoring (SAPS) are inadequate and not validated to predict which patients are likely to benefit from intensive care. [8-11]. Various pediatric triage system has been evaluated and analysed its association with the following surrogate clinical outcome measures of severity: hospitalisation rate, intensive care unit (ICU) admission, length of ED stay, predictive value for admission and length of hospitalization [12-16].

The most common being Pediatric Assessment Triangle (PAT) which is a rapid evaluation tool that establishes a child's

clinical status and his or her category of illness to direct initial management priorities [17]. PAT can be relied as only objective early warning of children in or at high risk for clinical deterioration but does not define PICU admission. All these triage system requires modifications targeted to young children and children with a comorbid conditions and sometimes misclassifies a substantial number of children who require ICU admission [18].

In addition to physiologic parameters and diagnoses, interpretation of the context of illness (acute vs exacerbation of chronic vs worsening of terminal illness), social implications, and religious beliefs may also be taken into consideration when determining admission to the PICU. Lastly, local socioeconomic context and limitation of healthcare resources must be considered the application of PICU admission criteria.

Pediatric critical care units in India face many challenges. In the government sector of the health system, there are few critical care units that are well equipped and that have the expertise to use sophisticated life sustaining technology. Furthermore, pediatric intensive care is poor or non-existent at district hospitals in rural India, where 80% of the nation's population resides and overcrowding of PICUs in urban settings is common [18-20]. Currently there is a lack of universally accepted, peer-reviewed recommendations for PICU admission criteria resource-limited settings and, in India national standards for pediatric critical care admission, practice and quality of care measures have not been established. Efficient use of intensive care services from a health resource standpoint is critical for several reasons. First, because intensive care is a precious commodity, especially in resource-limited settings, clarity about criteria for PICU admission assists local governments with resource allocation and service provision planning. Second, accurate categorization of patients in the emergency department setting shortens the time it takes to admit critically ill children to the proper care environment and also reduces unnecessary admissions for those who could be cared for safely and appropriately in a lower intensity setting. Lastly, standardized PICU admission criteria may be adopted and integrated by clinical personnel, hospitals, and health administrators to create local, regional, and national PICU care standards in context of location, environment and available resources. The current lack of recommendations is associated with significant provider variation in identifying pediatric intensive care needs and inconsistent use of PICU resources [21]. Once standard protocols and standardized indications of PICU admission are developed, India will move toward a more cost-effective use of its limited PICU resources [20]. Standardization of PICU admission criteria has been accomplished in developed countries through reviewed publications by professional societies [22], but its lacking in India. The purpose of this manuscript is to provide India specific recommendations which can be adapted to the local context and integrated into routine medical practices through a designated clinical and administrative body.

## Purpose and Intended Application

The purpose of these recommendations is to provide a framework and reference for future policy development by professional societies and governments in India. These recommendations are intended as a consensus outline, but should be adapted to meet the operational needs of each institution they are applied in, depending on the scope of illnesses encountered and the resources available. The definition of medical necessity for PICU admission reaches beyond India and general concepts outlined here may be utilized across resource-limited environments in different meetings. Application of these recommendations beyond the Indian context is feasible and suggestions for a process of implementation, monitoring, and evaluation are also included. Once health policies have been created, policy compliance along with clinical and administrative outcomes should be monitored by health administrators designated to oversee PICU care in institutions. Pediatric intensive care policies should be reviewed on a regular basis and revised as needed based on available evidence to support change.

## Consensus Recommendations Development Process

### Consensus panel task force

The consensus process applied is based on a previous approach by the Society of Critical Care Medicine [21], defining PICU admission criteria in high resource environments. These consensus recommendations were developed by a consensus panel task force team comprised of Indian and international experts in pediatric critical care, emergency medicine, trauma, and health policy stakeholders. Members were identified during the Indo-US Emergency and Trauma Collaborative conference 2015 (INDUSEM - Delhi) as leaders in intensive care policies from a variety of backgrounds in India and internationally.

These individuals were invited to participate in a discussion and consensus meeting during the 2016 annual INDUSEM WORLD CONGRESS at Bengaluru, India (Annex 1: Task force team members). In preparation for the 2016 consensus meeting, a consensus panel task force steering committee completed a global literature search about PICU admission criteria development, reviewed PICU recommendations published by a variety of professional organizations worldwide, and performed a literature review of relevant publications (Annex 2: Publications reviewed). The task force core group (Annex 1) performed a Pubmed literature search using Mesh Terms [intensive care] [pediatrics] [admission criteria] and identified relevant peer reviewed publications. In addition the group reviewed previously published statements from professional societies in India and other LMIC and compiled relevant publications in a literature resource list consisting of

400+ publications (Annex 2). The literature resource list was shared with the remaining consensus team members while the core group developed an initial draft of an evidence based list of conditions potentially relevant for PICU admissions in the resource-limited context of India [22-28]. Furthermore, based on previous approaches, the steering committee developed a framework for discussion and review of potential PICU parameters and defined the target outputs for the consensus meeting [29,30].

### Consensus process

The entire consensus panel task force team was assembled for an in-person round table discussion at the Indo-US Emergency and Trauma Collaborative conference during the 2016 INDUSEM WORLD CONGRESS in Bengaluru, India. Team members reviewed and discussed the various PICU admission criteria that were identified during the previous literature review and presented by members of the core group at the consensus meeting. The expert consensus panel then discussed and ranked proposed criteria according to scientific evidence, current standard of care, and expert opinion. Review to recommendation process: Based on field of practice, scientific expertise and location of practice we assemble subgroup teams (consensus panel core group members) who can provide content, specialty, research and methodological expertise in the review process and who were the primary drivers in drafting evidence based reviews and recommendations which were then further discussed by the full task force team until final consensus was obtained.

### Rating and decision making models

The decision about the necessity and appropriateness of PICU care was based on a variety or a combination of factors. Our consensus team followed a previously utilized approach to determine need of ICU admission based on a) prioritization modeling; b) general clinical criteria; c) clinical and objective parameters and d) other criteria [22,24].

**Levels of recommendation:** During the consensus process, meeting members applied following previously validated recommendation rating system [21].

**Level 1:** PICU admission justifiable on scientific evidence alone.

**Level 2:** PICU admission reasonably justifiable on scientific evidence and strongly supported by consensus expert opinion.

**Level 3:** Scientific evidence generally lacking but supported by available data and critical care expert opinion.

### Consensus panel task force recommendations on criteria for PICU admission

Recommendations on location of pediatric intensive care provision – High Dependency Units (Table 1).

**Table 1** Pediatric Intensive Care Provision in High Dependency Units

Pediatric Intensive Care Provision In High Dependency Units	Level of Recommendation
Pediatric intensive care can be provided at various locations within a healthcare facility. In addition to a designated PICU, many hospitals within India operate a High Dependency Unit (HDU) where intensive care can be provided, however staffing ratios and available equipment standards may differ from a standard PICU setup. The consensus task force panel identifies conditions which may be eligible to be cared for in a HDU setting if medical care for a specific condition can be delivered with equal quality when compared to the PICU setting. Conditions identified as eligible for HDU care are marked with an asterisk (*).	2
The minimum care standard for the HDU includes:  Minimal staffing requirements: 1:3 nurse to patient ratio; 1 resident level provider is available 24/7 to provide optimal medical supervision. The resident should be trained in pediatric advanced life support skills. The nurse should have substantial pediatric expertise.	3
Minimum services available to all patients: continuous cardiorespiratory monitoring; oxygen, suction, continuous monitoring, non-invasive ventilation modality, crash cart, defibrillator, lab 24/7, arterial blood gas, portable x-ray.  HDU must have immediate access to a dedicated PICU within their facility or have a relationship with an institution that has a PICU which can readily accept transfers if a patient can no longer be safely be managed in a HDU setting.	3
The minimum care standard for the PICU includes:  Unit design, equipment, organization and staffing and ancillary support services as recommended by ISCCM and IAP [25].	3

Recommendations on prioritization criteria for patients considered for PICU admission (Table 2). benefit most=Priority 1--to those who will benefit the least=Priority 4.

Assigning appropriateness for PICU admission based on a rating system which defined the patient populations who will

**Table 2** Prioritization Model Based PICU Admission

Prioritization model based PICU admission	Level
Priority 1:  Critically ill, unstable patients.  Patients who require monitoring, lifesaving or life sustaining treatment that cannot be provided outside the PICU Extent and duration of therapy are not limited by preexisting conditions or patient/family wishes  Examples  Respiratory failure requiring ventilator support Continuous vasoactive drug infusions (pressors, milrinone) Acute decompensated shock with signs of end organ failure Intentional or unintentional drug overdose, poisoning with end organ failure.	1
Priority 2:  Patients who require intensive monitoring and MAY need lifesaving or life sustaining treatment in near future  Examples  Severe respiratory distress with impending respiratory failure requiring possibly ventilator support. Shock responded to fluid boluses and MAY require monitoring for need of pressors.	1
Priority 3:  Critically ill patients with underlying life limiting illness  Limits in place as to extent of therapy (i.e patients with co-morbid conditions whose parents or guardians have decided against receiving resuscitation and /or lifesaving interventions)  Examples  Metastatic malignancy complicated by infections.	1
Priority 4:  PICU admission is not indicated  Monitoring and care can be provided outside PICU setting  Examples  Respiratory illnesses without evidence of active or impending respiratory failure.	1

## General clinical conditions that warrant PICU admission

Ideally, a patient should be admitted to the PICU setting before the condition reaches a point from where recovery is not possible. The minimum standards of PICU regarding the unit design, equipment, and organization and staffing as described. Early identification of clinical warning signs is important and requires health personnel who are trained and equipped to perform cardio-respiratory and neurologic assessments/interventions and to have decision-making skills. If a patient is diagnosed with a critical illness at a healthcare facility which does not have the capacity to provide the appropriate level of care, transfer to a higher level facility should be initiated immediately after the patient has been stabilized to the greatest extent possible.

General clinical conditions and indications warranting PICU admission

- All respiratory or cardiac arrest,

**Table 3** Clinical diagnosis model based PICU admission criteria

Clinical diagnosis model based PICU admission criteria	Level of recommendation
<b>Cardiac conditions</b>	
Cardiogenic shock, myocardial dysfunction: infectious and other	1
Complex dysrhythmias requiring close monitoring and intervention, including new onset complete heart block and after cardioversion	1
Acute congestive heart failure requiring hemodynamic support	1
Hypertensive emergencies	1
After cardiac arrest and post-resuscitation	1
Aortic dissection	1
Congenital heart disease with cardiopulmonary instability	1
Patients presenting to the emergency department with cardiorespiratory or neurologic compromise after high risk intrathoracic or cardiac procedures	1
Need for invasive cardiac monitoring	1
Need for cardiac pacing	1
Pericardial effusion requiring drainage, signs of tamponade	1
Hypertensive urgency	3*
<b>Pulmonary conditions</b>	
Acute respiratory insufficiency or failure requiring invasive mechanical ventilation	1
Hemoptysis with shock or airway compromise	1
Newborns with signs of severe respiratory distress	1
Rapidly progressive upper or lower respiratory disease with risk of progression to respiratory failure	1
High supplemental oxygen need >6 lpm or non-rebreather mask or FiO <sub>2</sub> >50% on CPAP/BiPAP to keep oxygen>94%	1*
Acute barotraumas (i.e decompression illness)	1*

- Unstable airway,
- Inability to oxygenate (O<sub>2</sub> Sat less than 90% on >50% oxygen requirement,
- Inability to ventilate with rising PCO<sub>2</sub> levels with respiratory insufficiency,
- Glasgow Coma Scale (GCS) score <8 or sudden fall in score by >2 points,
- Status epilepticus,
- Critical values of age specific vital signs parameters.

Clinical diagnosis and objective parameters that warrant PICU admission

This model uses specific well-defined clinical conditions which warrant PICU admissions (Table 3).

Numeric labels 1-3 designate level of recommendations (see above).

Asterisk indicates that such conditions can potentially be managed in an HDU.

Asthma-need for continuous administration of inhaled or nebulized medications to prevent respiratory failure	1*
Risk of complete airway obstruction	1
BRUE (brief resolved unexplained event) – recurrent	2*
<b>Neurologic conditions</b>	
Status epilepticus which cannot be controlled well with more than 2 antiepileptic medications (different class)	1*
Progressive neuromuscular dysfunction with altered mental status (GCS < 8 or <10 and deteriorating), respiratory or cardiovascular compromise	1
Non-traumatic Intracranial hemorrhage with evidence of increased ICP	1
Acute non-traumatic intracranial hemorrhage (epidural, subdural, subarachnoid, parenchymal)	1
Chronic progressive CNS disorders with deteriorating neurologic or respiratory function	1
Spinal cord compression or acute spinal lesions	1
Stroke with acute presentation	1*
Neurosurgical procedures requiring invasive monitoring of ICP	1
Hypertensive encephalopathy with PRES changes on imaging	1
Glasgow coma scale: GCS<8 → ICU; 9-13 → ICU or HDU	1, 1*
<b>Toxicologic conditions</b>	
Ingestions leading to severe neurologic compromise (GCS<8 or<10 and deteriorating) or respiratory compromise	1
Ingestions known to be associated with a high risk or cardio-respiratory events (e.g. recent organophosphate poisoning)	1*
Ingestions leading to hemodynamic instability, bleeding or organ failure.	1
Seizures following drug ingestion	1
Envenomation (snake/scorpion/bee stings)	1
<b>Gastrointestinal disorders</b>	
GI bleeding leading to hemodynamic instability, altered mental status or acidosis	1
Esophageal perforation	1
After emergency removal of foreign bodies	1*
Hepatic encephalopathy Grade>2	1
Corrosive ingestion	1
<b>Endocrinologic conditions</b>	
Diabetic keto-acidosis with hemodynamic instability, altered mental status, respiratory insufficiency or severe acidosis (pH<7.1)	1
Diabetic keto-acidosis with severe acidosis (pH<7.1) but without hemodynamic instability, altered mental status, or respiratory insufficiency	1
Hyperosmolar state with altered mental status and or hemodynamic instability	1
Adrenal crisis with hemodynamic instability	1
Inborn errors of metabolism with risk of respiratory, cardiovascular or neurologic decompensation	1*
Thyroid storm with hemodynamic instability	1
<b>Surgical or post-surgical conditions presenting in the emergency department setting</b>	
Patients after recent surgery presenting with hemodynamic, neurologic or respiratory compromise	1
Patient with a recent history of congenital heart disease repair presenting with hemodynamic, neurologic or respiratory compromise	1

Patients with recent open-intrathoracic surgeries presenting with hemodynamic, neurologic or respiratory compromise	1
Patients with recent organ transplantation presenting with hemodynamic, neurologic or respiratory compromise	1
<b>Radiologic findings</b>	
Cerebral vascular hemorrhage of any type with mental status change or focal neurologic signs	1
Ruptured viscera, bladder, uterus, liver esophagus	1
Bleeding of any type with hemodynamic instability	1
Dissecting aortic aneurisms	1
Foreign body before extraction with risk of perforation: batteries, sharp	2*
Tension pneumothorax	1*
Pleural effusion with cardiovascular or respiratory compromise	1*
Mediastinal mass with risk of obstruction	1*
Pulmonary embolism on CT<5d	1
<b>Children with special conditions – malignancies and hematologic conditions</b>	
Exchange transfusions	1
Plasmapheresis or leukopheresis	1*
Severe coagulopathy with active or high risk of bleeding	1
Severe complications of sickle cell diseases such as acute chest syndrome, aplastic anemia or hemodynamic instability	1
Tumor lysis syndrome	1*
Tumors or masses threatening airway, vital vessels or organs	1*
Febrile neutropenia with airway and hemodynamic compromise	1
<b>Conditions associated with trauma</b>	
Multiple trauma injury	1
Head trauma with acutely increased ICP, ANY evidence of cerebral edema on imaging	1
Severe head injury with altered mental status, respiratory compromise	1
Traumatic brain injury with GCS<8 or<10 and deteriorating	1
Traumatic brain injury in patient with bleeding disorder or receiving anti-coagulation therapy	1
Cardiac contusion, pulmonary contusion	1
Patients requiring placement of an extra ventricular drainage device (EVD)	1
Acute spinal cord injury	1
Trauma with intraabdominal organ injury	1*
Flail chest	1
Pelvic fracture with retroperitoneal hematoma	1
Crush injury	1
Grade 3 or 4 solid organ injury	1
<b>Burns</b> (regardless of underlying etiology)	Per ATLS recommendations
Trauma + 1 of the following	1

Requires massive blood transfusion, Base deficit>5, Seizures, Pregnancy, Hypothermia, Co-morbid conditions	
Placement recommendation Patients with severe traumatic injuries, intraabdominal injuries, TBI, GCS<8, crush injuries, or those likely requiring urgent surgical interventions should preferentially be admitted to ICU with availability of pediatric surgery and neurosurgery	
Intensive pain care needed: PCA, initiation of continuous infusion of opiates	1*
<b>Objective parameters, laboratory parameters</b>	
Potassium>6+clinical symptoms (with arrhythmias or weakness) Potassium>6 without clinical symptoms with or without EKG changes	1 2*
Potassium<2.5+clinical symptoms (with arrhythmias or weakness)	1
Ca>4 or iCa>10+/-clinical symptoms (hemodynamic instability or altered mental status (GCS<8 or<10 and deteriorating)	1
Ca 12-14 or iCa 8-10+clinical symptoms	2*
Ca<8 with or without symptoms (e.g. seizures)	1*
Hyponatremia with Serum Na<125 mmol/l or hypernatremia>160 mmol/l with clinical symptoms (e.g. altered mental status or seizures)	1*
Hyponatremia with Na<125 mmol/l without symptoms	3*
HgB<5+symptoms	1*
HgB<7 with active bleeding	1
<b>Other conditions</b>	
Shock of any etiology	1
Invasive Hemodynamic monitoring	1
Services not available at lower level care center: staffing shortages, drug shortages, equipment shortages	1*
Renal failure and need for acute hemodialysis	1*
Crush injury with acute renal insufficiency	1
Documented or suspected malignant hyperthermia	1
Snakebites and insect bites associated with cardiopulmonary or neurologic compromise as defined in respective sections	2*

## Administrative Recommendations to Facilitate Appropriate Paediatric ICU Admission

This document is designed to serve as a resource for hospitals and policy makers in resource-limited settings to determine appropriateness of PICU admissions for optimal utilization of available scarce resources within their own care environment.

Local stakeholders must take steps to achieve integration of PICU admission criteria into hospital care standards and health. Recommendations must be interpreted and applied in the local context of care, resources and health policy and should be adapted to meet the local needs. For successful integration into clinical practice, a hospital or region must appoint a physician director on the basis of qualification and leadership skill. This individual must be able to provide clinical, administrative, and educational direction to local staff to

integrate these recommendations into standard medical practice. Quality improvement processes need to be implemented to assure patient safety, to monitor compliance and to appropriate steps for continuous refinement of local policies.

Collaboration and integration of nursing staff, ancillary staff, and directors of other units within the hospital is essential to ensure transparency of the quality improvement process. The ultimate decision responsibility for acceptance and refusal of PICU admission is in the hands of the transferring and accepting physician, who may deviate from the recommendation if this deviation is in the best interest of the patient. Ideally a multidisciplinary team should conduct non-threatening reviews of protocol deviations, adverse patient events, and hospitalization outcomes in order to further refine applicability of these recommendations. By establishing a culture that focuses on systems issues and re-education as opposed to blame and punishment, institutions will find it

more feasible to be in compliance with best practice standards, where care is safe, effective, and efficient.

## Limitations of applicability of these recommendations

Even though every effort was made to identify all relevant literature, it is possible that important publications may have been missed in the search. Some references used date back to the 1980's indicating the paucity of available literature in this topic especially with application on low resource settings such as India. Due to the complexity of medical conditions under review, high variability in the quantity and quality of literature covering the spectrum of medicine and ICU indications, our team decided to utilize the level 1-3 rating system [19] over more traditional Evidence level A-E rating system.

Even though every effort was made to have reputable experts in emergency medicine, pediatrics and intensive care with a variety of medical and working backgrounds participate in the consensus process, it may be possible that some practitioners may have been over- and some underrepresented. Even though literature review and drafting evidence based recommendations for final review and inputs was accomplished by team members with topic specific clinical, research and methodological research, we did not include subspecialists in the consensus process. Due to the complexity of health care systems within India between the public and private sector, variation in staffing, staff competency, availability of equipment between hospitals, urban-rural healthcare delivery discrepancies, state and institution specific variable definitions of HDU, staffing standards, etc. and considering a variety of other factors, the authors realize that a uniform application of these recommendations is not possible and is also not intended. The authors see this publication as a reference and starting point for institutions who are interested in engaging in the process of defining PICU admission criteria. These recommendations are also not designed or intended to serve as ethical or medico-legal criteria to be applied to decide about "appropriateness" of care, placement of patients and transfer of patients and are not meant to replace clinical judgment and the local definition of appropriate care. Overcrowding, high caseloads exceeding hospital capacity and limited bed availability in HDU and PICU units are commonly encountered in India; however these recommendations are not designed to address eligibility of transfer-in and transfer-out policies in these units and provide a universally applicable recommendation on overflow scenarios.

## Summary

This publication is designed to provide recommendation of clinical criteria for PICU admissions for children from the emergency department. The authors see this publication as a reference and starting point for institutions who are interested in engaging in the process of defining PICU admission criteria, to assist key stakeholders in the development of hospital operational standards, to define appropriateness of PICU

admission will assist in effective resource allocation, maximize healthcare benefits for the population, reduce healthcare resource waste, and improve access to quality care for children. This publication discusses clinical conditions and scenarios that warrant PICU or HDU admission but is not intended to be utilized as an ethical or medical-legal document but as a resource for clinicians, hospital and systems administrators to standardize care processes, reduce variation in care. Recommendations are provided based on prioritization modeling as well as on clinical conditions.

## Acknowledgements

We would like to acknowledge the INDUSEM organization, which was the driving force behind the development of these recommendations. We would further like to acknowledge the Government of India Ministry of Health and Family Welfare, Medical Council of India, Indian Academy of Pediatrics, Indian Society of Critical Care Medicine and Shakti Krupa Charitable Trust for their support of this project.

## References

1. Bone RC, McElwee NE, Eubanks DH (1993) Analysis of indications for intensive care unit admission-Clinical efficacy project-American College of Chest Physicians. *Chest* 104: 1806-1811.
2. NIH Consensus Conference-Critical Care Medicine (1983) *JAMA* 2506: 798-804.
3. Mulley AG (1983) The allocation of resources for medical intensive care. In: President's commission for the study of ethical problems in medicine and biomedical research: Securing access to health care. Washington DC, Government Printing Office, 3: 285-311
4. Kollef MH, Shuster DP (1994) Predicting ICU outcomes with scoring systems: Underlying concepts and principles. *Crit Care Clin* 10: 1-18.
5. Nasraway SA, Cohen IL, Dennis RC (1998) Recommendations on admission and discharge for adult intermediate care units. *Crit Care Med* 26: 607-610.
6. Charlson ME, Sax FL (1987) The therapeutic efficacy of critical care units from two perspectives: A traditional cohort approach vs a new case-control methodology. *J Chron Dis* 40: 31-39.
7. Ron A, Aronne LJ, Kalb PE (1989) The therapeutic efficacy of critical care units: Identifying subgroups of patients who benefit. *Arch Intern Med* 149: 338-341.
8. Pollack MM, Ruttimann E, Getson PR (1988) Pediatric risk of mortality (PRISM) score. *Crit Care Med* 16: 1110-1116.
9. Smith G, Nielsen M (1999) ABC of intensive care criteria for admission. *BMJ* 218: 1544-1547.
10. Stenson MD (1992) Scarcity in the intensive care unit: Principles of justice for rationing ICU beds. *Am J Med* 92: 552-555.
11. Wagner DP, Knaus WA, Harrell FE (1994) Daily prognostic estimates for critically ill adults in intensive care units: Results from a prospective, multicenter, inception cohort analysis. *Crit Care Med* 22: 1359-1372.

12. Gravel J, Fitzpatrick E, Gouin S (2013) Performance of the Canadian triage and acuity scale for children: A multicenter database study. *Ann Emerg Med* 61: 27-32.e3.
13. Gravel J, Manzano S, Arsenault M (2009) Validity of the Canadian Paediatric Triage and Acuity Scale in a tertiary care hospital. *Crit J Emerg Med* 11: 23-28.
14. Roukema J, Steyerberg EW, van Meurs A (2006) Validity of the Manchester Triage System in paediatric emergency care. *Emerg Med J* 23: 906-910.
15. Green NA, Durani Y, Brecher D (2012) Emergency Severity Index version 4: A valid and reliable tool in pediatric emergency department triage. *Pediatr Emerg Care* 28: 753-757.
16. Ganapathy S, Yeo JG, Thia XHM, Hei GMA, Tham LP, et al. (2018) The Singapore Paediatric Triage Scale Validation Study. *Singapore Med J* 59: 205-209.
17. Horeczko T, Enriquez B, McGrath NE, Gausche-Hill M, Lewis RJ, et al. (2013) The Pediatric Assessment Triangle: accuracy of its application by nurses in the triage of children. *J Emerg Nurs* 39: 182-189.
18. Vidyasagar D, Singh M, Bhakoo O (1997) Evolution of Neonatal and Pediatric Critical Care in India. *Critical Care Clinics* 13: 331-347.
19. Yeolekar ME, Mehta S (2008) ICU care in India – Status and Challenges. Editorial – *J Associat Phys India* 56: 221-222.
20. Udwadia F, Guntupalli K (1997) Critical care in India. *Crit Care Clin* 13: 317-320.
21. Prayag S (2002) ICUs worldwide: Critical care in India. *Crit Care* 6: 479-480.
22. Nates J, Nunnally M, Kleinpell R (2016) ICU Admission, Discharge and Triage recommendations: A Framework to enhance clinical operations, development of institutional policies and further research. *Crit Care Med* 44: 1553-1602.
23. Clinical Practice Recommendations for Admission to the Pediatric Intensive Care Unit (2010) MOH Paediatrics Clinical Recommendations. CWMH & Lautoka Hospital.
24. Smith G, Nielsen M (1999) ABC of intensive care criteria for admission. *Bri Med J* 318: 1544-1547.
25. AAP Policy Statement: American Academy of Pediatrics-Committee on Hospital Care and Section on Critical Care (1999) Recommendations for developing admission and discharge policies for the pediatric intensive care unit. *Pediatrics* 103: 840-842.
26. Recommendations for Intensive Care Unit Admission, Discharge and Triage (1997) Taskforce of the American college of critical care medicine, Society of Critical Care Medicine. *Crit Care Med* 27: 633-638.
27. Rosenberg D, Moss M, The American College of Critical Care Medicine of the Society of Critical Care Medicine (2004) Recommendations and levels of care for pediatric intensive care units. *Crit Care Med* 32: 2117-2127.
28. Khilnani P (2002) Indian Society of Critical Care Medicine (Pediatric Section); Indian Academy of Pediatrics (Intensive Care Chapter). *Indian Pediatr* 39: 43-50.
29. Govil YC (2006) Pediatric intensive care in India: Time for introspection and intensification. *Indian Pediatr* 43: 675-867.
30. Vidyasagar D, Singh M, Bhakoo ON (1997) Evolution of neonatal and pediatric critical care in India. *Crit Care Clin* 13: 331-346.