Phytoremediation: An eco-friendly approach for recovery of soils contaminated by heavy metals

Antonio Cristaldia, Gea Oliveri Contia*, Salvatore Luciano Cosentinob, Giovanni Mauromicaleb, Chiara Copata, Alfina Grassoa, Pietro Zuccarelloa, Maria Fiorea, Cristina Restucciab, Margherita Ferrantea

aEnvironmental and Food Laboratories (LIAA), Department of Medical Science, Surgical and Advanced Technologies "G.F. Ingrassia", University of Catania, Italy.
bDepartment of Agriculture, Food and Environment (Di3A), University of Catania, Italy.

Corresponding Author: Gea Oliveri Conti

✉ antonio.cristaldi@unict.it

Citation: Cristaldia A, The Known Sub-Sequence Algorithm (KSSA) for making optimal decisions about imputation methods for multiple missing data in time series for environmental research. Glob Environ Health Saf.2022, 6:S1:4

Abstract

Missing data in time series is a frequent problem for the environmental sciences. This is a serious limitation for statistical analysis and therefore, imputation (the process of filling missing data) is a keystone task. Several imputation methods have been proposed and implemented in programming software, however, their efficiency is data-dependent. There is no universal imputation method best for all-time series, but instead, each method suits the structure of particular groups of time series. ¿Which imputation method is best to fill a time series? the main problem is that the target time series (of interest for imputation) already contains missing data, so validation of methods cannot be performed directly on it. Instead it needs a full time series (no missing data) to simulate missing data, perform imputations and compare actual to impute. However, the best imputation method for the full time series is not necessarily the best for the target time series. The Known Sub- Sequence Algorithm (KSSA) is a novel approach to solve this problem by validating imputation methods directly on target time series. It uses the information contained within sub-sequences between missing data gaps to produce an optimal decision about a best imputation method for any particular target time series, no matter the structure it has. This is done by means of a process of iterative bootstrapping that randomly samples sub-sequences of the target time series in order to learn from them to find a best method form a set of candidates. This is a promising machine learning algorithm that will help environmental scientists and decision makers working with time series. KSSA will soon be implemented as the 'kssa' R-package in CRAN and is currently available in GitHub.

Biography

Ivan Felipe Benavides is Biologist from Universidad de Nariño in Colombia, PhD in Ecology from Universidad Austral de Chile, current postdoc researcher at Universidad Nacional de Colombia, and consultant for SoftwareShop and Datambiente in Latin America for environmental issues. He is an expert in environmental data science, which includes biostatistics, modeling, experimental design, machine learning and algorithm development to solve environmental problems.

References

1. Ali, A., Guo, D., Mahar, A., Wang, P., Shen, F., Li, R., Zhang, Z., 2017. Mycoremediation of Potentially Toxic Trace Elements – a Biological Tool for Soil Cleanup: A Review. Pedosphere 27(2), 205–222.[CrossRef] [GoogleScholar] [Indexed at]

2. Ashraf, S., Ali, Q., Zahir, Z.A., Ashraf, S., Asghar, H.N., 2019. Phytoremediation: Environmentally sustainable way for reclamation of heavy metal polluted soils. Ecotoxicology and Environmental Safety 174, 714–727.[CrossRef] [GoogleScholar] [Indexed at]

3. Cristaldi, A., Oliveri Conti, G., Jho, E.H., Zuccarello, P., Grasso, A., Copat, C., Ferrante, M., 2017. Phytoremediation of contaminated soils by heavy metals and PAHs. A brief review. Environmental Technology & Innovation 8, 309–326.[CrossRef] [GoogleScholar] [Indexed at]

4. Fiorentino, N., Fagnano, M., Adamo, P., Impagliazzo, A., Mori, M., Pepe, O., Ventorino, V., Zoina, A., 2013. Assisted phytoextraction of heavy metals: compost and Trichoderma effects on giant reed (Arundo donax L.) uptake and soil N-cycle microflora. Ital. J. Agron. 8 (4), 244–254.[CrossRef] [GoogleScholar] [Indexed at]

5. Cristaldi, A., Oliveri Conti, G., Cosentino, S.L., Mauromicale, G., Copat, C., Grasso, A., Zuccarello, P., Fiore, M., Restuccia, C., Ferrante, M., 2020. Phytoremediation potential of *Arundo donax* (Giant Reed) in contaminated soil by heavy metals. Environmental Research, 185, 2020, 109427.[CrossRef] [GoogleScholar] [Indexed at]