

# Prospects of Biological Control Agents for Management of Invasive Pest *Tuta Absoluta* (Lepidoptera: Gelechiidae) In Africa

Never Zekeya<sup>1\*</sup>, Humphrey Peter<sup>2</sup>, Ernest Mbega<sup>3</sup>

<sup>1</sup>Department of Animal sciences, College of African Wildlife Management-Mweka, Moshi, Tanzania

<sup>2</sup>Department of Animal sciences, Tanzania Industrial Research and Development Organization, Msasani, Dar es Salaam, Tanzania

<sup>3</sup>Department of Sustainable Agriculture and Biodiversity Ecosystem Management, School of Life Science and Bioengineering, The Nelson Mandela African Institution of Science and Technology, Arusha, Tanzania

\*Corresponding Authors: Never Zekeya, Department of Animal sciences, College of African Wildlife Management-Mweka, Moshi, Tanzania, E-Mail: Wairimuann012@gmail.com

Received date: December 03,2021; Accepted date: December 16,2021 ; Published date: December 23,2021

Citation: Zekeya N, Peter H, Mbega E (2021) Prospects of Biological Control Agents for Management of Invasive Pest *Tuta Absoluta* (Lepidoptera: Gelechiidae) In Africa. J Anim Sci Livest Prod Vol:5 No:6

## Abstract

*Tuta absoluta* (Meyrick) is a devastating agricultural pest of global concern, which has spread widely across continents causing substantial economic losses, particularly in tomato cultivation. In Africa the problem is much severe to the extent that farmers are abandoning the crop. Controlling *T. absoluta* by use of chemical pesticides is profoundly tricky due to development of rapid resistance. Alternative options such as biological controls are either scarce or expensive and only a handful of microbial agents are currently registered for management of *T. absoluta* in Africa. It has been observed that *T. absoluta* causes damage and loss in tomato ranging from 50-100% in South America, Europe and Africa where management options are most limited. Although several synthetic pesticides are registered for management of *T. absoluta* including; chlorantraniliprole spinetoram and emamectin benzoate group, little success are reported in Africa due to various factors including fast resistance development of *T. absoluta*. Commercially available biopesticides such as; *Beauveria bassiana* (Bals.), *Metarhizium anisopliae* (Metschn.), *Bacillus thuringiensis* (Bt), neem extract (Azadirachtin) and *Spinosad* (bacteria compound; spinosyn A and spinosyn D) are scarcely distributed in Africa. Although recently there are newly studied biocontrols including *Aspergillus oryzae* (Vuruga Biocide) that is registered in Tanzania, yet not distributed to most farmers across the country. However, some

biocontrols have been reported for control *T. absoluta* in Africa including; *Aspergillus flavus* (Ahlb.) and *Aspergillus oryzae* (Ahlb.), they are not commercially available. This review highlights on the utilization and commercialization of various biological control agents for *T. absoluta* management in Africa.

**Keywords:** Tomato leafminer; *Aspergillus oryzae*; Management strategies; Microbial control agents

## Introduction

*Tuta absoluta* (Meyrick), the tomato leafminer, is an invasive pest from South America that has spread extensively in Europe, Asia and Africa threatening tomato production [1,2,3,4,5].

*T. absoluta* and other pests, including mealybug and white fly, aphids, cabbage looper, cutworms, leafminer (*Liriomyza sativa*), tomato fruit worm, and leaf beetles are the main threats to tomato producers in Africa and globally except USA where *T. absoluta* has not yet been properly reported [6,7,3,8,9,10,11].

*T. absoluta* and other lepidopterous pests reduce both the yield and the quality of tomato fruit, thus diminishing its value in the market [12,13,14].

**Table 1:** Global distribution of *Tuta absoluta*, host range and damage incidences.

Continent	Confirmed countries	Host	Damage	Invasion period	Reference
South America	Peru, Chile, Ecuador, Argentina, Brazil, Bolivia, Colombia, Paraguay, Uruguay, Venezuela, Peru, and	Tomato, potato, Datura spp, and Nicotiana spp	Damage range from 80-100% Damage range 80-100% tomato damage	1914 - to date; still persisting	Sylla et al., 2019; Desneux et al., 2011; [121] Luna et al., 2015

Central/ America	North	Canada, Haiti, USA, Mexico, Panama, Costa Rica	Tomato, potato, Datura spp, and Nicotiana spp	damage 30-50%	Present with moderate impact	Desneux et al., 2011; [125,126,128]
Europe		Spain, Italy, Malta, France, Portugal, Germany, Bulgaria, UK, Serbia, Switzaland, Slovenia, Romania, Lithuania, Greece, Cosovo, Azerbaijan, Croatia, Belarus, Netherlands, Belgium, Czech-Republic, Denmark, Georgia, Guernsey, Hungray, North Macedonia, Norway, ukraine, Austria, Montenegro, Cyprus, Bosnia and Herzegovina	Tomato, potato, and Datura spp, Nicotiana spp, Cape gooseberry, bean (Phaseolas vulgaris)	Damage rate range from 50-80% crop damage in protected cultivation	Late 2006 – to date. Continuing to invade new areas and cause damage	[111]; Karadjova et al., 2013; [124]
Asia		Turkey, India, Iran, Japan, China, Rusia, Yemen, Qatar, Bahrain, Kuwait, United Arab Emirate, Bangladesh, Iraq, Syria, Sultanate of Oman, Lebanon, Israel, Palestine, Saudi Arabia, China, Nepal, Afghanistan, Kazakhstan, Tajikistan, Uzbekistan, Pakistan and Burma	Tomato and eggplant	50-100% tomato damage	2010 – onwards; gradually expand its range in the region	Li et al., 2021; [115] Guimapi et al., 2020; Zhang et al., 2020; Han et al., 2019; [106]; [119] Rakha et al., 2017; [131]
Africa		Algeria, Morocco, Libya, Egypt, Senegal, Sudan, Canary island, Eritrea, Nigeria, Ethiopia, Lesotho, Angola, Burundi, Cape Verde, Ghana, Botswana, Benin, Mayotte, Equatorial Guinea, Kenya, Tunisia, Niger, Tanzania, Zambia, Zimbabwe, D R Congo, Somalia, Bukina faso, Namibia, DR Sao Tome and Principe, Togo and Côte d'Ivoire	Tomato, potato, night shade, eggplant, Datura spp, and Nicotiana spp	Up to 100% crop damage in open fields	Since 2008 and currently expanding its invasion across the continent	[120,113,134] Sawadogo et al., 2020; Smith et al., 2018; [109] Desneux et al., 2011; Rakha et al., 2017; Tonnang et al., 2015

Australia	Not reported	None reported	Under risk due to high commercial	Expect invasion on tomato	Han et al., 2018; Verheggen et al., 2019
-----------	--------------	---------------	-----------------------------------	---------------------------	--

*Tuta absoluta* has spread globally and is now distributed across various agroecological regions of Africa causing losses tomato as well as other crops such as night shade, eggplant, pepper, and several weeds [15,16] (Table 1). The Tomato leaf miner is also a major impediment in the production of several other solanaceous vegetables in Africa [17,18], It has led to severe damage and yield losses, mainly due to insecticide resistance [19]. Since its arrival in 2014, Tanzanian farmers have experienced massive crop losses [20]. Its control has become a major challenge due to the broad host range and a higher reproductive rate [21,22]. *Tuta absoluta* attacks all developmental crop stages, from seedlings to mature fruits for field and greenhouse-grown solanaceas including *S. tuberosum*, *S. nigrum*, *S. melongena* and wild *Solanum spp* [23-25]. Studies have shown that *Tuta absoluta* larvae and adults prefer the aerial plant parts due to their higher sugar levels [26,27]. This pest is successful due to tolerance under extreme temperature conditions ranging from coldness to high temperature hence surviving in several regions and various seasons [28,29]. Hence, management of this pest is tricky requiring an effective, integrated and affordable biocontrol agent to avert the problem in Africa.

Management of *T. absoluta* in Tanzania and in other African countries rely heavily on application of conventional pesticides [30] without even detection of the pest by using pheromone and colored traps that assist in identification and monitoring of pest population [31]. Despite the popularity of chemical pesticides, studies have revealed the usefulness of using other options including host resistance mechanisms [32]. Generally, there are scarce reports on management options with ability of controlling all stages of *Tuta absoluta*. Biological pesticides including viruses, fungi, bacteria and nematodes have been reported to effectively control *T. absoluta* eggs and larvae [33,34]. However, few of these have undergone screening in Africa despite its widespread invasion.

Natural enemies, particularly Hemipteran predators, are effective against *T. absoluta* and other insect pests [35,36,37]. In addition, entomopathogenic microbes are also gaining popularity in insect pest management worldwide, including Africa [38,39,10,40].

The alternate use of different active ingredients biocontrol agents such as; predators, parasitoids, botanicals and microbials showed high efficacy with low or no rate resistance on the pest [8,41]. However, use of an integrated pest management including use of botanicals such as neem extract, microbial pesticide and minimal chemical pesticides is highly recommended for effective synegetic effects and safety with natural enemies [42,43,35]. Microbial controls including fungi, bacteria and virus have high potential to be used as biocontrol agents for management of insect pests for crop protection [33,44]. Bacteria especially *Bacillus spp* have proven to be effective for controlling *Tuta absoluta* in many areas [45]. Several reports have proved the

efficacy of entomopathogenic bacteria; *B. thuringensis* for the control of *T. absoluta* larvae in Africa [40,46]. However, the effectiveness and accessibility of these biopesticides vary with pest species and geographical areas whereas little success has been reported in controlling *T. absoluta* in Tanzania [47]. Entomopathogenic fungi have been highly used as foliar sprays/drenches directly on plant or as endophytes for controlling insect pests compared to the rest of microbial controls [48,49]. *Beauveria bassiana* is widely used as epiphyte or endophyte for pest control worldwide [48,50] with high efficacy as both an endophyte and a foliar spray. Other entomopathogenic fungi including *Metarhizium anisopliae* have been reported to be highly effective against different pests including *T. absoluta* [51]. On other hand, *Aspergillus spp.* have been screened and evaluated in laboratory with high efficacy against larvae, pupae and adults of *T. absoluta* in Algeria and Tanzania [52,47]. In Africa, entomopathogenic microbes have been introduced for pest management with few record of registered products [53].

Since the management of insect pests is shifting from reliance on chemicals to integrated approaches of biocontrols including pathogenic nematodes, bacteria, fungus and virus [54,55,56,53], it is evident that biopesticides will increasingly be adopted in many regions especially in Africa where the pest is recently invading abruptly. Hence, native microbial controls including *Aspergillus* species could be a potential biopesticide for control of *T. absoluta* as it has been revealed with promising results on *T. absoluta* in Africa [52,47]. This review highlights the global trends of *T. absoluta*, available and registered management options for controlling *T. Absoluta*. It also provides prospects for screening and registration of novel biopesticide from local context for effective management of *T. absoluta* in Africa where the pest available management options are unaffordable by most farmers.

## Current management strategies for controlling *Tuta absoluta*

### Conventional pesticides

Synthetic chemical pesticides have been used worldwide with low success due to resistance in most regions [51,57]. Management of *T. absoluta* by single approach cannot suppress the pest below economic injury level due to the fact that there is no single option that can 100% control all stages [58]. Several studies reported the failure of chemical pesticides including Chlorantraniliprole and Amamectin benzoate for controlling *T. absoluta* [19,59]. Chemical pesticides are imported in Africa in which their effectiveness might be compromised during shipping and misused by farmers due pest panic as well as limited training on application of pesticides [47]. Additionally, synthetic pesticides are environmentally and economically unaffordable in most African countries with limited financial resources. However, proper use with integration with environmentally safe

strategies including traps and biological agents should be promoted to enhance management of *T. absoluta*.

#### Pheromone traps

Pheromones and other semiochemicals are widely applied for controlling insect pests in integrated pest management (IPM) programs including monitoring, mass trapping, lure and killing (attract-inhale) and mating disruption [32]. Studies show that pheromone traps are effective in controlling *T. absoluta* under protected cultivation [60]. However, although traps are useful for attracting mainly adult male moth and often for population monitoring [5], they are expensive, and are rarely used by farmers to control *T. absoluta* in Africa. Studies show that pheromone traps cannot be effective when used merely for control of pests because they are designed to disrupt mating and not killing [61]. However, other studies have revealed that incorporation of semiochemicals; attractants of different colors in pheromone traps in early growing season help detect pests and monitor population and gradually minimize pest impact in the field [32,5]. Proper selection and application of traps could be effective when integrated with biological controls to safely protect tomato and other vegetable crops against *T. absoluta*.

#### Microbial control agents against *T. absoluta*

Microbial controls for pest management are becoming popular worldwide as effective and affordable control options against various insect pests on various crops [38]. In Europe and America, the application of microbial controls for crop protection has been commercialized and used intensively [62] whereas it has only been recently introduced in Africa. Microbes play a major role in balancing nutrients and proper utilization in

agriculture [63]. A study by Buragohain et al. 2021 reported the effectiveness of integrating *B. thuringiensis var kurstaki*, *B. bassiana* and neem extract for management of *T. absoluta* under field conditions in India. This reveals that native biocontrol agents are potential for effective management of various insect pests [64,65,66]. Other microbial controls including viruses, bacteria and nematodes have been reported to be effective against eggs and larvae of *T. absoluta* under laboratory conditions [33,34] but their further validation is needed. Despite the fact that *T. absoluta* is widespread in Africa, few microbial controls have been registered and commercialized [47,67]. The commercially available bacterial biopesticides are *B. thuringiensis* and *B. sub illis* [55,68,65,69,70]. Recently, entomopathogenic fungi are increasingly studied and reported to be effective pest controls [71,72,73]. Some fungi species including *B. bassiana* are reported as plant endophytes enhancing pest control [48,6], and also produce toxic metabolites for pest control [74,75]. On the other hand, *M. anisopliae*, *B. bassiana*, *A. latus* and *A. clavatus* have shown high efficacy against several agricultural insect pests [76]. It has also been observed that most entomopathogenic fungi are effective in controlling lower stages of pests and some crop diseases when combined with chemical pesticides [77]. However, improving access, training on application and handling of available biopesticides particularly *M. anisopliae*, *B. bassiana* and *B. thuringiensis* and proper integration with other pesticides are critical, which could ensure potential market in Tanzania and Africa at large [78]. Since microbial control options for insect pest management are becoming popular globally, it is high time to embark on their commercialization in Africa.

**Table 2:** Current management strategies for controlling *Tuta absoluta* in invaded regions.

Type	Commercial name/ ingredients & source	Efficacy	Accessibility	Reference
Synthetic pesticides	Chlorantraniliprole (Coragen, E.I. DuPont de Nemours and Company, New York, USA); Emamectin benzoate (Wiltigo, King Quenson Group, Shenzhen, China); spinetoram (Radiant, Dow Agrosiences, Indianapolis, USA); S	For controlling, eggs, larvae and pupae, but with high pest resistance within short time	Low application in America, Europe, Asia, recently introduced and misused in Africa with little success	[107,114]; Guedes & Picanço, 2012; [121,129]
Traps	Pheromone traps ( <i>Tuta absoluta</i> -Optima), Russell IPM Company (London, UK)	Contain 0.5 mg lure (TDTA and TDDA)/trap for trapping adults	America, Europe, Asia, Africa (limited access) mainly for monitoring	Desneux et al., 2010; [122] Ostrauskas & Ivinskis, 2010
Botanicals	<i>Azadirachta indica</i>	Eggs and larvae	Africa, America, Asia	Kona et al., 2014
Microbial control	<i>Metarhizium anisopliae</i> , <i>Beauveria bassiana</i> , <i>Bacillus thuringiensis</i> (Bt)	Larvae, eggs, pupae	Canada, USA, Australia, Spain, Mexico, Colombia, South Africa,	Fite et al., 2020; [102]; Contreras et al., 2014; Sabbour & Nayera, 2014; [130]

	<i>Aspergillus oryzae</i> ( <i>Vuruga Biocide</i> )		Kenya, Egypt, and Tanzania	
Nematodes	<i>Steinernema carpocapsae</i> , <i>Steinernema feltiae</i> , <i>Heterorhabditis bacteriophora</i>	Larvae, pupa and adults	No commercialization information, no access in developing countries	Batalla-Carrera et al., 2010
Predator	<i>Nabis pseudoperous</i> <i>Blattostethus pallescens</i> <i>Poppius la</i>	Larvae, larvae and eggs (under laboratory)	Spain	[104,117]
Parasitoids	<i>Trichogramma achaeae</i> , <i>T. pretiosum</i> , <i>Neochrysocharis formosa</i> , <i>Necremnus tutae</i> , <i>Necremnus near artynes</i> and <i>Necremnus near tidius</i>	Eggs and larvae	Europe and America, Argentina	de Campos et al., 2020; Ballal et al., 2016; Biondi et al., 2013; [108]; Ferracini et al., 2012; Luna et al., 2011; Pratissoli & Parra, 2000
Resistant varieties tomato	<i>Unappealing leaf surface</i> , <i>trichome</i> , <i>repellant cues</i>	<i>Egg, larvae, adult repellant</i>	None are commercially adopted	[123] Rakha et al., 2017; [132]
Protected cultivation	<i>Application of green/net houses</i>	<i>Preventing pest attack physically</i>	America, Europe, Asia but less adopted in African countries	Cocco, Deliperi, & Delrio, 2013; [112]; Ostrauskas & Ivinskis, 2010

### Microbial pesticide potential for commercialization

*Aspergillus* species particularly *A. oryzae* have been revealed to infect several insect life stages [47] registered (BCA/IN0005) for controlling *Spodoptera frugiperda* (fall armyworm) and *T. absoluta* in Tanzania. This novel microbial control agent could further be evaluated in other African counties as biocontrol of *T. absoluta* and other pests. Other fungal species such as

*Aspergillus* species have been reported to control various pests in tropical countries [52,79], hence should be further evaluated and commercialized for control of *T. absoluta* in Africa. This review informs researchers to screen and commercialize new microbial control agents including *A. oryzae* that is reported to control different stages of *T. absoluta* including adult females to effective control crop damage in the field (Table 3).

**Table 3:** List of potential microbial control agents for controlling *T. absoluta*.

Name	Target	Active ingredient/utility	Application	Reference
<i>Bacillus thuringensis</i> , <i>Bacillus subtilis</i>	Eggs, larvae	Spore/colony forming units commercially available in America, Europe, Asia and Africa	Foliar sprays and seed coats for seed and seedling protection	Abdullah, 2020; Kamal et al., 2019; Gowtham et al., 2018; Sabbour, 2014; Abd El-Ghany et al. 2016b
<i>Metarhizium anisopliae</i>	Eggs, larvae	Conidia/metabolites commercialized in America and Europe	Applied as direct foliar spray, root endophytes of conidia	[127] Contreras et al., 2014)
<i>Beauveria bassiana</i>	Eggs, larvae and pupae	Conidial formulations commercialized in developed countries	Endophytes and foliar spray of formulated conidia	[116], Inanli, Yoldaş, & Birgücü, 2012; Urbaneja et al., 2012; Abd El-Ghany et al. 2016b
<i>Trichoderma album</i> , <i>Trichoderma harzianum</i>	Eggs, larvae and adults	Conidia of <i>T. album</i> and <i>T. harzianum</i> in a neem	Laboratory tested but not commercialized	Kaoud, 2014; [118,135-137]

			blend no commercial product for <i>T. absoluta</i>		
<i>Aspergillus flavus</i> , <i>Aspergillus oryzae</i> (BCA/IN/0005)	Eggs, larvae, pupae and adult	Infective conidia of <i>A. flavus</i> reported in laboratory where <i>A. oryzae</i> - registered in Tanzania for controlling lepidopterous pests	With broad application as biopesticide and potential endophyte of crops in Africa	[110,138,139]; Lakhdari et al., 2016; Zekeya et al., 2019; FAO-highlights, 2020	

### Natural enemies

Parasitoids and predators including *Macrolophus pygmaeus* and Hymenopteran, respectively, are effective in controlling *T. absoluta* eggs and larvae and several insect pests helping to reduce damage and suppress population [80-83]. Table 2 shows a list of parasitoids with their revealed efficacy in developed countries [27] few parasitoids are reported for controlling *T. absoluta* in Africa [80]. However, lack of expertise on parasitoids rearing and parasitism efficacy could be one of the obstacles of utilizing this approach for controlling insect pests in Africa. Hence there is a need for screening and registration of parasitoids as natural controls of *T. absoluta* in the region.

### Botanical extracts


Botanical extracts have been and are currently used for controlling insect pest although only few exist commercially [84,85]. Neem extract has been widely used for the management of crop pests with efficacy against *T. absoluta* eggs, larvae and pupae [1]. Another study by [103] revealed the potential of using piper extracts for control of *T. absoluta* on tomato. Studies revealed that botanical extracts are potential sources of active ingredients for controlling pests, particularly *T. absoluta* and can have enhanced activity when combined with synthetic pesticides [86]. A study by [105] revealed that Orange peel oil have high insecticidal activity against larvae of *T. absoluta*. Again, other studies revealed that plant extracts are potential larvicides of *T. absoluta* when prepared and applied appropriately. However, botanical pesticides should be careful handled as they might have side effects as synthetic chemical pesticides in some circumstances [87-89]. Hence, this review highlights potential biological pesticides due to high efficacy, potential for commercialization and safety for sustainable crop production.

### Future outlook for controlling *Tuta absoluta* in Africa

Integrated pest management approach could be effective for sustainable control of *Tuta absoluta* [90]. Biopesticides from *B. bassiana*, *M. anisoplia*, *B. subtilis* and *B. thuringiensis* have promising results for controlling a broad range of pests [38,91,66]. However, their effectiveness on host pests is diminutive, due to context variations compared to native strains that have shown high efficiency on *T. absoluta* [66].

Thus, there is an urgent need for screening for effective alternative options available in local contexts. Figure 1 summarizes the potential applicability of genus *Aspergillus* as a biocontrol agent and shows how *Aspergillus* species can

promote plant growth through production of growth hormones such as gibberellins [63,92].



**Figure 1:** The potential for formulation of novel biopesticide and various modes of application of *Aspergillus* species as endophyte/epiphyte enriching plant nutrient levels (NL) in soil upon saprophytic activity and pest control via foliar application to kill eggs, larvae, pupae and moth suppressing pest below economic injury level (EEL) in fields.

It also shows that *Aspergillus spp* produces phytohormones and has insecticidal activities against agricultural pests [76] and can act as bio-fertilizers by decomposing organic materials [101]. *Aspergillus* species has a good record in nutrient enhancement and promotion of healthy plant growth [93] whereas *A. flavus* and *A. awamori* act upon industrial waste water and heavy metals, providing nutrients and minerals and crop protection [94-96]. *A. fumigatus* and *A. parasiticus* are reported to have endophytic activity in plants through production of essential minerals and metabolites [97]; *A. oryzae* has been reported as a marine endophyte that produces protective metabolites against pests [98]. *Aspergillus* species are also known for producing metabolites that enhance plant defense against pests and diseases (Sun et al., 2018) and act as a source of metabolites in various environments [99]. Furthermore, [79] have reported activity of *A. oryzae* against locusts and, more recently, *A. flavus* has shown significant activity against *T. absoluta* in Algeria [52]. Locally available entomopathogenic fungi could be potential suppressors of all life stages of *T. absoluta* including adult moths, and reduce its impact in Africa [47]. However, integrated pest management and biological control are essential for enhancing efficacy [100]. Hence an *Aspergillus*-based biopesticide could have multiple and integrated applications with other biological control agents. It could be incorporated in irrigation systems, foliar sprays and drenching applications to maximize pest control efforts

## Conclusion

*Tuta absoluta* is a tricky pest to be controlled by a single strategy. Various management options are available in other regions but with little success in Africa. Besides a wider use of synthetic pesticides for management of *T. absoluta* in most regions, their effectiveness is context based due to varied agroecological conditions leading to failures and increasing cultivation cost. Due to these shortcomings of the available pest management options, this review recommends for commercialization of locally available biocontrol agents in Africa. However, screening for novel microbial agents should be funded and promoted to enhance efficacy, accessibility and applicability for effective management of destructive pest, *T. absoluta* in Africa.

## References

- Konan K A J, Ouali-Ngoran S W M, Fondio L, Ochou G O, Koné D, et al. (2021) Geographical distribution and host range status of *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) in Côte d'Ivoire. *Int J Trop Insect Sci* 1-13
- Guimapi R A, Srinivasan R, Tonnang H E, Sotelo-Cardona P, & Mohamed S A (2020) Exploring the Mechanisms of the Spatiotemporal Invasion of *Tuta absoluta* in Asia. *Agriculture* 10:124
- Han P, Bayram Y, Shaltiel-Harpaz L, Sohrabi F, Saji A, et al. (2019) *Tuta absoluta* continues to disperse in Asia: damage, ongoing management and future challenges. *J Pest Sci* 92:1317-1327
- Berxolli A, & Shahini S (2017) Population dynamic of tomato leaf miner, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae). *J agric sci* 85-89
- Abd El-Ghany N M, Abdel-Wahab E S, & Ibrahim S S (2016a) Population fluctuation and evaluation the efficacy of pheromone-based traps with different color on tomato leafminer moth, *Tuta absoluta* (Lepidoptera: Gelechiidae) in Egypt. *Res J Pharm Biol Chem* 7:1533-1539
- Wei Q Y, Li Y Y, Xu C, Wu Y X, Zhang Y R, et al. (2020) Endophytic colonization by *Beauveria bassiana* increases the resistance of tomatoes against *Bemisia tabaci*. *Arthropod-Plant Interactions* 1-12
- Verheggen F and Fontus R B (2019) First record of *Tuta absoluta* in Haiti. *Entomologia Generalis* 38(4):349-353
- Mansour R, & Biondi A (2020) Releasing natural enemies and applying microbial and botanical pesticides for managing *Tuta absoluta* in the MENA region. *Phytoparasitica* 1-16
- Rakha M, Zekeya N, Sevgan S, Musembi M, Ramasamy S, et al. (2017) Screening recently identified whitefly/spider mite-resistant wild tomato accessions for resistance to *Tuta absoluta*. *Plant Breeding* 136:562-568
- Abd El-Ghany N M, Abdel-Razek A S, Djelouah K, & Moussa A (2018) Efficacy of bio-rational insecticides against *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) on tomatoes. *Bios Res* 15:28-40
- Ibrahim SS, Moharum FA, Abd El-Ghany NM (2015) The cotton mealybug *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae) as a new insect pest on tomato plants in Egypt. *J Plant Prot Res* 55:48-51
- Dhaliwal G, Vikas J, & Dhawan A (2010) Insect pest problems and crop losses: changing trends. *Indian J Ecol* 37:1-7
- Harizanova V, Stoeva A, & Mohamedova M (2009) Tomato leaf miner, *Tuta absoluta* (Povolny) (Lepidoptera: Gelechiidae)—first record in Bulgaria. *J Agric Sci Technol* 1:95-98
- Kutinkova H, Caicedo F, & Lingren B (2016) The main pests on Solanacea crops in Zona 1 of Ecuador. *New Know J of Sci* 5
- Idriss G E, du Plessis H, Khamis F M, Ekesi S, Tanga C M, et al. (2020) Host Range and Effects of Plant Species on Preference and Fitness of *Tuta absoluta* (Lepidoptera: Gelechiidae). *J Econ Entomol*
- Abbes K, Harbi A, Elimem M, Hafsi A, & Chermiti B (2016) Bioassay of three solanaceous weeds as alternative hosts for the invasive tomato leafminer *Tuta absoluta* (Lepidoptera: Gelechiidae) and insights on their carryover potential. *African Entomology* 24:334-342
- Sylla S, Brévault T, Monticelli L S, Diarra K, & Desneux N (2019) Geographic variation of host preference by the invasive tomato leaf miner *Tuta absoluta*: implications for host range expansion. *J Pest Sci* 92:1387-1396
- Tonnang H E, Mohamed S F, Khamis F, & Ekesi S (2015) Identification and risk assessment for worldwide invasion and spread of *Tuta absoluta* with a focus on Sub-Saharan Africa: implications for phytosanitary measures and management. *PLoS ONE* 10:0135283
- Campos MR, Silva T B, Silva W M, Silva J E, & Siqueira H A (2015) Spinosyn resistance in the tomato borer *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *J Pest Sci* 88:405-412
- Materu C L, Shao E A, Losujaki E, Chidege M, & Mwambela N (2016) Farmer's perception knowledge and practices on management of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in tomato growing areas in Tanzania. *Int J Res* 1
- Maerere P A, Mwatawala MW, Mtui H D, Mamiro D P, & Baltazari A (2020) Temporal distribution and management of the tomato leaf miner. *Int J Veg Sci* 1-11
- Smith J D, Dubois T, Mallogo R, Njau EF, Tua S, et al. (2018) Host range of the invasive tomato pest *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) on solanaceous crops and weeds in Tanzania. *Florida Entomologist* 573-579
- Silva G A, Queiroz EA, Arcanjo L P, Lopes M C, Araújo T A, et al. (2021) Biological performance and oviposition preference of tomato pinworm *Tuta absoluta* when offered a range of Solanaceous host plants. *Scientific Reports* 11:1-10
- Zekeya N, Ouma J, Chacha M, Ndossi H, & Mbega E R (2020) *Bios Res* 17:874-880
- Karadjova O, Ilieva Z, Krumov V, Petrova E, & Ventsislavov V (2013) *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae): Potential for entry, establishment and spread in Bulgaria. *Bulg J Agric Sci* 19:563-571
- Terzidis A N, Wilcockson S, & Leifert C (2014) The tomato leaf miner (*Tuta absoluta*): Conventional pest problem, organic management solutions? *Organic Agriculture* 4: 43-61
- De Campos M R, Monticelli L S, Bearez P, Amiens-Desneux E, Wang Y, et al. (2020) Impact of a shared sugar food source on biological control of *Tuta absoluta* by the parasitoid *Necremnus tutae*. *J Pest Sci* 93:207-218

28. De Campos M R, Béarez P, Amiens-Desneux E, Ponti L, Gutierrez AP, et al. (2021) Thermal biology of *Tuta absoluta*: demographic parameters and facultative diapause. *J Pest Sci* 94:829-842
29. Li X W, Li D, Zhang Z J, Huang J, Zhang J M, et al. (2021) Supercooling capacity and cold tolerance of the South American tomato pinworm, *Tuta absoluta*, a newly invaded pest in China. *J Pest Sci* 94:845-858
30. Sawadogo M W, Somda I, Nacro S, Legrève A, & Verheggen F J (2020) Insecticide susceptibility level and control failure likelihood estimation of Sub-Saharan African populations of tomato leafminer: Evidence from Burkina Faso. *Physiological Entomology* 45:147-153
31. Heidari N, Sedaratian-Jahromi A, Ghane-Jahromi M, & Zalucki M P (2020) How bottom-up effects of different tomato cultivars affect population responses of *Tuta absoluta* (Lep.: Gelechiidae): a case study on host plant resistance. *Arthropod-Plant Interactions* 14:181-192
32. Abd El-Ghany NM (2019) Semiochemicals for controlling insect pests. *J Plant Prot Res* 59:1-11
33. Abd El-Ghany NM, Abdel-Razek AS, Mahmoud YA, Ebadah IMA (2016b) Evaluation of some microbial agents, natural and chemical compounds for controlling tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *J Plant Prot Res* 56:372-379
34. Luna M G, Pereyra P C, Coviella C E, Nieves E, Savino V, et al. (2015) Potential of biological control agents against *Tuta absoluta* (Lepidoptera: Gelechiidae): current knowledge in Argentina. *Florida Entomologist* 98:489-494
35. Mansour R, Brévault T, Chailleux A, Cherif A, Grissa-Lebdi K, et al. (2018) Occurrence, biology, natural enemies and management of *Tuta absoluta* in Africa. *Entomol Gen* 38:83-112
36. Zappala L, Biondi A, Alma A, Al-Jboory I J, Arno J, et al. (2013) Natural enemies of the South American moth, *Tuta absoluta*, in Europe, North Africa and Middle East, and their potential use in pest control strategies. *J Pest Sci* 86:635-647
37. Al-Jboory IJ, Katbeh-Bader A, & Al-Zaidi S (2012) First observation and identification of some natural enemies collected from heavily infested tomato by *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae) in Jordan. *Middle East J Sci Res* 11:435-438
38. Fite T, Tefera T, Negeri M, Damte T, & Sori W (2020) Evaluation of *Beauveria bassiana*, *Metarhizium anisopliae*, and *Bacillus thuringiensis* for the management of *Helicoverpa armigera* (Hubner)(Lepidoptera: Noctuidae) under laboratory and field conditions. *Biocontrol Sci Technol* 30:278-295
39. Abdel-Razek AS, Abd El-Ghany NM, Djelouah K, Moussa A (2017) An evaluation of some eco-friendly biopesticides against *Bemisia tabaci* on two greenhouse tomato varieties in Egypt. *J Plant Prot Res* 57:9-17
40. Abd El-Ghany NM, Abdel Ghany EM, Salama HS (2015a) Efficiency of *B. thuringiensis* Isolates from Egypt against the Pink Bollworm *Pectinophora gossypiella* (Saunders). *Biopesticide International* 11:12- 19
41. Ghoneim K (2014) Predatory insects and arachnids as potential biological control agents against the invasive tomato leafminer, *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae), in perspective and prospective. *J Entomol Zool Stud* 2:52-71
42. Buragohain P, Saikia D K, Sotelo-Cardona P, & Srinivasan R (2021) Development and validation of an integrated pest management strategy against the invasive South American tomato leaf miner, *Tuta absoluta* in South India. *Crop Protection* 139: 105348
43. Akutse K S, Subramanian S, Khamis F M, Ekesi S, & Mohamed S A (2020) Entomopathogenic fungus isolates for adult *Tuta absoluta* (Lepidoptera: Gelechiidae) management and their compatibility with *Tuta* pheromone. *J Appl Entomol*
44. Biondi A, Chailleux A, Lambion J, Han P, Zappalà L, et al. (2013) Indigenous natural enemies attacking *Tuta absoluta* (Lepidoptera: Gelechiidae) in southern France. *Egypt J Biol Pest Control* 23:117
45. Gowtham V, Kannan M, Senthilkumar M, & Soundararajan R P (2018) Isolation and characterization of indigenous *Bacillus thuringiensis* Berliner from animal ordure effective against South American tomato pinworm, *Tuta absoluta* (Meyrick). *Pest Management in Horticultural Ecosystems* 24:1-7
46. Sabbour M (2014) Biocontrol of the tomato pinworm *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. *MEJAR* 3:499-503
47. Zekeya N, Mtambo M, Ramasamy S, Chacha M, Ndakidemi PA, et al. (2019) First record of an entomopathogenic fungus of tomato leafminer, *Tuta absoluta* (Meyrick) in Tanzania. *Biocontrol Sci Technol* 29:626-637
48. Nishi O, Sushida H, Higashi Y, & Iida Y (2020) Epiphytic and endophytic colonisation of tomato plants by the entomopathogenic fungus *Beauveria bassiana* strain GHA. *Mycology* 1-9
49. Ekesi S, Dimbi S, & Maniania N (2007) The role of entomopathogenic fungi in the integrated management of fruit flies (Diptera: Tephritidae) with emphasis on species occurring in Africa. *Research SignPost: Kerala, India* 37:239-274
50. Meyling N V, & Eilenberg J (2007) Ecology of the entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae* in temperate agroecosystems: potential for conservation biological control. *Biological Control* 43:145-155
51. Biondi A, Guedes RNC, Wan F-H, & Desneux N (2018) Ecology, worldwide spread, and management of the invasive South American tomato pinworm, *Tuta absoluta*: past, present, and future. *Annu Rev Entomol* 63:239-258
52. Lakhdari W, Dehliz A, Acheuk F, Mlik R, Hammi H, et al. (2016) Biocontrol test against the leaf miner of tomato *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) by using entomopathogenic fungi in the Algerian Sahara. *J algér rég Arid* 13:85-89
53. Kamali S, Karimi J, & Koppenhöfer A M (2018) New insight into the management of the tomato leaf miner, *Tuta absoluta* (Lepidoptera: Gelechiidae) with entomopathogenic nematodes. *J Econ Entomol* 111:112-119
54. Guedes R, & Picanço M (2012) The tomato borer *Tuta absoluta* in South America: pest status, management and insecticide resistance. *EPPO Bulletin* 42:211-216
55. Kaoud H (2014) Alternative methods for the control of *Tuta absoluta*. *Global Impact Factor* 2:41-46
56. Sabbour M, & Nayera Y S (2014) Evaluations of three *Bacillus thuringiensis* against *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. *Int. J Sci Res* 3:2319-7064
57. Passos L C, Soares M A, Collares L J, Malagoli I, Desneux N, et al. (2018) Lethal, sublethal and transgenerational effects of insecticides on *Macrolophus basicornis*, predator of *Tuta absoluta*. *Entomol Gen* 38:127-143

58. Khani S, Hejazi M J, & Karimzadeh R (2020) Susceptibility of eggs and larvae of tomato leafminer, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) to some insecticides. *JESI* 39:393-402
59. Khan H A A, Akram W, & Ali S (2020) Activities of Select Enzymes Involved in Insecticide Resistance in Spinosad-Resistant and-Susceptible Strains of *Musca domestica* (Diptera: Muscidae). *J Med Entomol* 57:620-622
60. Jallow M F, Dahab A A, Albaho M S, Devi V Y, Jacob J, et al. (2020) Efficacy of mating disruption compared with chemical insecticides for controlling *Tuta absoluta* (Lepidoptera: Gelechiidae) in Kuwait. *Appl Entomol Zool* 1-9
61. Cocco A, Deliperi S, & Delrio G (2013) Control of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in greenhouse tomato crops using the mating disruption technique. *Appl Entomol* 137:16-28
62. Luna M G, Wada V I, La Salle J, & Sanchez N E (2011) *Neochrysocharis formosa* (Westwood) (Hymenoptera: Eulophidae), a newly recorded parasitoid of the tomato moth, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae), in Argentina. *Neotrop. Entomol* 40:412-414
63. Khan A L, Hamayun M, Kim Y-H, Kang SM, Lee J-H, et al. (2011) Gibberellins producing endophytic *Aspergillus fumigatus* sp. LH02 influenced endogenous phytohormonal levels, isoflavonoids production and plant growth in salinity stress. *Process Biochemistry* 46:440-447
64. Salama HS, Abd El-Ghany NM, Saker M (2015) Diversity of *Bacillus thuringiensis* isolates from Egyptian soils as shown by molecular characterization. *J Genet Eng Biotechnol* 13:101- 109
65. Salama HS, Saker M; Salama M, El-Banna A, Ragaei M et al. (2012) *Bacillus thuringiensis* isolates from Egyptian soils and their potential activity against lepidopterous insects. *Arch Phytopathol Plant Prot* 45:856-868
66. Merdan A, Salama HS, Labib E, Ragaei M and Abd El-Ghany NM (2010) *Bacillus thuringiensis* Isolates from Soil and Diseased Insects in Egyptian Cotton Fields and Their Activity Against Lepidopterous Insects. *Arch Phytopathol Plant Prot* 43:1165 - 1176
67. Barra-Bucarei L, Moreno LD, & Iglesias A F (2019) Biological Control in Chile. *Biological Control in Latin America and the Caribbean: Its Rich History and Bright Future* 12:108
68. Saker M, Salama HS, Ragaei M, and Abd El-Ghany NM (2012) Molecular Characterization of *Bacillus thuringiensis* Isolates from The Egyptian Soil. *Arch Phytopathol Plant Prot* 45:110-125
69. González-Cabrera J, Mollá O, Montón H, & Urbaneja A (2011) Efficacy of *Bacillus thuringiensis* (Berliner) in controlling the tomato borer, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *BioControl* 56:71-80
70. Batalla-Carrera L, Morton A, & García-del-Pino F (2010) Efficacy of entomopathogenic nematodes against the tomato leafminer *Tuta absoluta* in laboratory and greenhouse conditions. *BioControl* 55:523-530
71. Kamal M I, El-Adl AM, Zaid K A, El-Sayed A I, & Atia H H (2019) Genetic Diversity among Transconjugants of *Bacillus thuringiensis* Used for Controlling Tomato Borer *Tuta absoluta*. *J Agric Food Chem* 10:231-236
72. Abd El-Ghany N M, Lotfy D E, & Embaby E M (2015b) Isolation and evaluation of native *Metarhizium anisopliae* strains against the greater wax moth *Galleria mellonella* Linnaeus (Lepidoptera: Pyralidae). *Egypt J Biol Pest Control* 199
73. Contreras J, Mendoza J, Martínez-Aguirre M, Garcia-Vidal L, Izquierdo J, & Bielza P (2014) Efficacy of entomopathogenic fungus *Metarhizium anisopliae* against *Tuta absoluta* (Lepidoptera: Gelechiidae). *J Econ Entomol* 107:121-124
74. Abdullah R R H (2020) The efficacy of Chitinolytic bacteria as a bioinsecticide against *Tuta absoluta* (Lepidoptera: Gelechiidae). *Int J Entomol Res* 5:107-110
75. Ishak I, Ng L C, Haris-Hussain M, Jalinas J, Idris A B, et al. (2020) Pathogenicity of an Indigenous Strain of the Entomopathogenic Fungus *Metarhizium anisopliae* (Hypocreales: Clavicipitaceae) (MET-GRA4 Strain) as a Potential Biological Control Agent Against the Red Palm Weevil (Coleoptera: Dryophthoridae). *J Econ Entomol* 113:43-49
76. Sun B T, Akutse K S, Xia XF, Chen J H, Ai X, et al. (2018) Endophytic effects of *Aspergillus oryzae* on radish (*Raphanus sativus*) and its herbivore, *Plutella xylostella* *Planta* 248:705-714
77. Dhanya M K, Sathyan T, Murugan M, Ashokkumar K, & Surya R (2019) In-vitro Compatibility of Entomopathogenic Fungi with Synthetic Insecticides and Neem Oil. *Pestic Res J* 31:275-281
78. Van Lenteren J C, Bolckmans K, Köhl J, Ravensberg W J & Urbaneja A (2018) Biological control using invertebrates and microorganisms: plenty of new opportunities. *BioControl* 63:39-59
79. Zhang P, You Y, Song Y, Wang Y, & Zhang L (2015) First record of *Aspergillus oryzae* (Eurotiales: Trichocomaceae) as an entomopathogenic fungus of the locust, *Locusta migratoria* (Orthoptera: Acrididae). *Biocontrol Sci Technol* 25:1285-1298
80. Aigbedion-Atalor P O, Mohamed S A, Hill M P, Zalucki M P, Azrag A G, et al. (2020) Host stage preference and performance of *Dolichogenidea gelechiidivoris* (Hymenoptera: Braconidae), a candidate for classical biological control of *Tuta absoluta* in Africa. *Biological Control* 144:104215
81. Gupta A (2020) Hymenopteran Parasitoids in Cultivated Ecosystems: Enhancing Efficiency. In *Innovative Pest Management Approaches for the 21st Century*. Springer, Singapore 323-338
82. Ferracini C, Ingegno B L, Navone P, Ferrari E, Mosti M, et al. (2012) Adaptation of indigenous larval parasitoids to *Tuta absoluta* (Lepidoptera: Gelechiidae) in Italy. *J Econ Entomol* 105:1311-1319
83. Ballal CR, Gupta A, Mohan M, Lalitha Y, & Verghese A (2016) The new invasive pest *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in India and its natural enemies along with evaluation of Trichogrammatids for its biological control. *Current science* 2155-2159
84. Ndereyimana A, Nyalala S, Murerwa P, & Gaidashova S (2020) Field efficacy of entomopathogens and plant extracts on *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) infesting tomato in Rwanda. *Crop Protection* 105183
85. Hafsi A, Abbes K, Chermiti B, & Nasraoui B (2012) Response of the tomato miner *Tuta absoluta* (Lepidoptera: Gelechiidae) to thirteen insecticides in semi-natural conditions in Tunisia. *EPPO bulletin* 42:312-316
86. Soares M A, Campos M R, Passos L C, Carvalho GA, Haro M M, et al. (2019a) Botanical insecticide and natural enemies: a potential combination for pest management against *Tuta absoluta*. *J Pest Sci* 92:1433-1443
87. Arnó J, & Gabarra R (2011) Side effects of selected insecticides on the *Tuta absoluta* (Lepidoptera: Gelechiidae) predators *Macrolophus pygmaeus* and *Nesidiocoris tenuis* (Hemiptera: Miridae). *J Pest Sci* 84:513-520
88. Ghoneim K (2014) Parasitic insects and mites as potential biocontrol agents for a devasative pest of tomato, *Tuta absoluta*

- Meyrick (Lepidoptera: Gelechiidae) in the world: a review. *IJRR* 19:1
89. Isman M B (2015) A renaissance for botanical insecticides?. *Pest management science* 71:1587-1590
  90. Maerere P A, Mwatawala M W, Mtui H D, Mamiro D P, & Baltazari A (2021) Temporal distribution and management of the tomato leaf miner. *Int J Veg Sci* 27:120-130
  91. Shalaby H, Faragalla F, El-Saadany H, & Ibrahim A (2013) Efficacy of three entomopathogenic agents for control the tomato borer, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Natural Science* 11:63-72
  92. Tsavkelova E, Klimova S Y, Cherdyntseva T, & Netrusov A (2006). Microbial producers of plant growth stimulators and their practical use: a review. *Appl Biochem Microbiol* 42:117-126
  93. Pérez-García, A, Romero D, & De Vicente A (2011) Plant protection and growth stimulation by microorganisms: biotechnological applications of Bacilli in agriculture. *Curr Opin Biotechnol* 22:187-193
  94. Mittal V, Singh O, Nayyar H, Kaur J, & Tewari R (2008) Stimulatory effect of phosphate-solubilizing fungal strains (*Aspergillus awamori* and *Penicillium citrinum*) on the yield of chickpea (*Cicer arietinum* L. cv. GPF2). *Soil Biol Biochem* 40:718-727
  95. Ghatnekar S D, Kaviani M F, Sharma SM, Ghatnekar S S, Ghatnekar G S, et al. (2010) Application of vermi-filter-based effluent treatment plant (pilot scale) for biomanagement of liquid effluents from the gelatine industry. *Dynamic Soil, Dynamic Plant* 4:83-88
  96. Iram S, Ahmad I, & Stuben D (2009) Analysis of mines and contaminated agricultural soil samples for fungal diversity and tolerance to heavy metals. *Pak J Bot* 41:885-895
  97. Kusari S, Pandey S P, & Spiteller M (2013) Untapped mutualistic paradigms linking host plant and endophytic fungal production of similar bioactive secondary metabolites. *Phytochemistry* 91:81-87
  98. Debbab A, Aly A H, & Proksch P (2011) Bioactive secondary metabolites from endophytes and associated marine derived fungi. *Fungal Diversity* 49:1-12
  99. Gurulingappa P, Sword G A, Murdoch G, & McGee P A (2010) Colonization of crop plants by fungal entomopathogens and their effects on two insect pests when in planta. *Biological Control* 55:34-41
  100. Chailleux A, Desneux N, Seguret J, Khanh H D T, Maignet P, et al. (2012) Assessing European egg parasitoids as a mean of controlling the invasive 1. South American tomato pinworm *Tuta absoluta*. *PLoS ONE* 7:48068
  101. Bhardwaj D, Ansari M W, Sahoo RK, & Tuteja N (2014) Biofertilizers function as key player in sustainable agriculture by improving soil fertility, plant tolerance and crop productivity. *Microbial Cell Factories* 13:66
  102. Borgi I, Dupuy J-W, Blibech I, Lapaillerie D, Lomenech A-M, et al. (2016) Hyper-proteolytic mutant of *Beauveria bassiana*, a new biological control agent against the tomato borer. *Agron Sustain Dev* 36:60
  103. Brito E F D, Baldin E L L, Silva R D C M, Ribeiro L D P, & Vendramim J D (2015) Bioactivity of Piper extracts on *Tuta absoluta* (Lepidoptera: Gelechiidae) in tomato. *Pesqui Agropecu Bras* 50:196-202
  104. Cabello T, Gallego J, Fernandez-Maldonado F, Soler A, Beltran D, et al. (2009) The damsel bug *Nabis pseudoferus* (Hem.: Nabidae) as a new biological control agent of the South American tomato pinworm, *Tuta absoluta* (Lep.: Gelechiidae), in tomato crops of Spain. *OIBC/WPRS Bulletin*, 49:219-223
  105. Campolo O, Cherif A, Ricupero M, Siscaro G, Grissa-Lebdi K, et al. (2017) Citrus peel essential oil nanoformulations to control the tomato borer, *Tuta absoluta*: chemical properties and biological activity. *Scientific reports* 7:1-10
  106. Campos MR, Rodrigues A R S, Silva W M, Silva TB M, Silva V RF, et al. (2014) Spinosad and the tomato borer *Tuta absoluta*: a bioinsecticide, an invasive pest threat, and high insecticide resistance. *PLoS ONE* 9:103235
  107. Campos M R, Biondi A, Adiga A, Guedes R N, & Desneux N (2017) From the Western Palaearctic region to beyond: *Tuta absoluta* 10 years after invading Europe. *J Pest Sci* 90:787-796
  108. Chailleux A, Bearez P, Pizzol J, Amiens-Desneux E, Ramirez-Romero R, et al. (2013) Potential for combined use of parasitoids and generalist predators for biological control of the key invasive tomato pest *Tuta absoluta*. *J Pest Sci Science* 86:533-541
  109. Chermiti B, Abbes K, Aoun M, Ben Othmane S, Ouhibi M, et al. (2009) First estimate of the damage of *Tuta absoluta* (Lepidoptera: Gelechiidae) and evaluation of the efficiency of sex pheromone traps in greenhouses of tomato crops in the Bekalta region, Tunisia. *Afr J Plant Sci* 3:49-52
  110. Dahliz A, Lakhdari W, Soud A, Hammi H, Boucekima H, et al. (2013) Complex of natural enemies and control methods of the exotic invasive pest *Tuta absoluta* (Lepidoptera: Gelechiidae) in Southern Algeria. *Crop Protection Management in Mediterranean Organic Agriculture* 48
  111. Desneux N, Wajnberg E, Wyckhuys K A, Burgio G, Arpaia S, et al. (2010) Biological invasion of European tomato crops by *Tuta absoluta*: ecology, geographic expansion and prospects for biological control. *J Pest Sci* 83:197-215
  112. Desneux N, Luna M G, Guillemaud T, & Urbaneja A (2011) The invasive South American tomato pinworm, *Tuta absoluta*, continues to spread in Afro-Eurasia and beyond: the new threat to tomato world production. *J Pest Sci* 84:403-408
  113. Fiaboe K R, Agboka K, Agboyi L K, Koffi D, Ofuo R, Kpadonou G E et al. (2020) First report and distribution of the South American tomato pinworm, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae) in Togo. *Phytoparasitica* 1-11
  114. Gontijo P, Picanço M, Pereira E, Martins J, Chediak M, & Guedes R (2013) Spatial and temporal variation in the control failure likelihood of the tomato leaf miner, *Tuta absoluta*. *Ann Appl Biol* 162:50-59
  115. Gui-fen Z H A N G, De-ying M A, Yu-sheng W A N G, You-hua GA O, Wan-xue L IU (2020) First report of the South American tomato leafminer, *Tuta absoluta* (Meyrick), in China. *J Integr Agric* 19:1912-1917
  116. Inanli C, Yoldaş Z, & Birgücü A (2012) Effects of entomopathogenic fungi, *Beauveria bassiana* (Bals.) and *Metarhizium anisopliae* (Metsch.) on larvae and egg stages of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Ege Üniversitesi Ziraat Fakültesi Dergisi* 49:239-242
  117. Jamwal R, Sharma P L, Verma S C, Chandel RS, & Sharma N (2021) Demographics and functional response of *Blaptostethus pallascens* preying on *Tuta absoluta*. *Phytoparasitica* 1-13
  118. Khidr A, Gaffar S, Maha S, Nada A, Taman A, et al. (2013) New approaches for controlling tomato leafminer, *Tuta absoluta* (Meyrick) in tomato fields in Egypt. *Egypt J Agric Res* 91:335-348

119. Kiliç T (2010) First record of *Tuta absoluta* in Turkey. *Phytoparasitica* 38:243-244
120. Kona N E M, Taha A K, & Mahmoud M E (2014) Effects of Botanical Extracts of Neem (*Azadirachta indica*) and *Jatropha* (*Jatropha curcus*) on eggs and larvae of tomato leaf miner, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae). *Persian Gulf Crop Protection* 3:41-46
121. Lietti M M, Botto E, & Alzogaray R A (2005) Insecticide resistance in argentine populations of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Neotrop Entomol* 34:113-119
122. Michereff Filho M, Vilela E F, Jham G N, Attygalle A, Svatos A, et al. (2000) Initial studies of mating disruption of the tomato moth, *Tuta absoluta* (Lepidoptera: Gelechiidae) using synthetic sex pheromone. *J Braz Chem Soc* 11:621-628
123. Oliveira C M D, Andrade Júnior V C D, Maluf W R, Neiva I P, & Maciel G M (2012) Resistance of tomato strains to the moth *Tuta absoluta* imparted by allelochemicals and trichome density. *Ciência e Agrotecnologia* 36:45-52
124. Ostrauskas H, & Ivinskis P (2010) Records of the tomato pinworm (*Tuta absoluta* (Meyrick, 1917))—Lepidoptera: Gelechiidae—in Lithuania. *Acta Zoologica Lituanica* 20:151-155
125. Parra J RP (2009) Egg parasitoids commercialization in the new world. In *Egg Parasitoids in Agroecosystems with Emphasis on Trichogramma* 373-388
126. Pereyra P C, & Sánchez N E (2006) Effect of two solanaceous plants on developmental and population parameters of the tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Neotrop Entomol* 35:671-676
127. Pires L M, Marques E J, Wanderley-Teixeira V, Teixeira Á A, Alves L C, et al. (2009) Ultrastructure of *Tuta absoluta* parasitized eggs and the reproductive potential of females after parasitism by *Metarhizium anisopliae*. *Micron* 40:255-261
128. Proffit M, Birgersson G, Bengtsson M, Reis R, Witzgall P, et al. (2011) Attraction and oviposition of *Tuta absoluta* females in response to tomato leaf volatiles. *J Chem Ecol* 37:565-574
129. Reyes M, Rocha K, Alarcón L, Siegwart M, & Sauphanor B (2012) Metabolic mechanisms involved in the resistance of field populations of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) to spinosad. *Pestic Biochem Physiol* 102:45-50
130. Sanahuja G, Banakar R, Twyman R M, Capell T, & Christou P (2011) *Bacillus thuringiensis*: a century of research, development and commercial applications. *Plant Biotechnol J* 9:283-300
131. Shashank P, Chandrashekar K, Meshram N M, & Sreedevi K (2015) Occurrence of *Tuta absoluta* (Lepidoptera: Gelechiidae) an invasive pest from India. *Indian j entomol* 77:323-329
132. Simmons A T, & Gurr G M (2005) Trichomes of *Lycopersicon* species and their hybrids: effects on pests and natural enemies. *Agric For Entomol* 7:265-276
133. Soares M A, Passos L C, Campos M R, Collares L J, Desneux N, et al. (2019b) Side effects of insecticides commonly used against *Tuta absoluta* on the predator *Macrolophus basicornis*. *J Pest Sci* 92:1447-1456
134. Zekeya N, Ndakidemi P A, Chacha M, & Mbega E (2017) Tomato leafminer, *Tuta absoluta* (Meyrick 1917), an emerging agricultural pest in Sub-Saharan Africa: Current and prospective management strategies. *Afr J Agric Res* 12:389-396
135. Fand B B, Shashank P R, Suroshe S S, Chandrashekar K, Meshram N M, et al. (2020) Invasion risk of the South American tomato pinworm *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae) in India: predictions based on MaxEnt ecological niche modelling. *Int J Trop Insect Sci* 1-11
136. Guevara E J, Porcel M, Calixto A M, Bueno V D P, & Moino A (2020) Interactions Between the Nematode *Heterorhabditis amazonensis* JPM4 and the Predator *Macrolophus basicornis*: Two Natural Enemies of *Tuta absoluta* Native to South America. *Neotrop Entomol* 49:108-115
137. Kumar B (2018) Insect Pest Management. In *Pests and Their Management* 1015-1078
138. Silva A C L, Silva G A, Abib P H N, Carolino A T, & Samuels R I (2020) Endophytic colonization of tomato plants by the entomopathogenic fungus *Beauveria bassiana* for controlling the South American tomato pinworm, *Tuta absoluta*. *CABI Agriculture and Bioscience* 1:1-9
139. Steiner M & Goodwin S (2015) *Tuta absoluta*. *Practical Hydroponics and Greenhouses* 153:38-44