

Pelagia Research Library

Asian Journal of Plant Science and Research, 2022, 12(4)



Response of Tef [Eragrostis tef (Zuccagni) Trotter] to Nitrogen and NPS-B Blended Fertilizers on Andisols of Southern Ethiopia

Melkamu Hordofa Sigaye*, Kidist Kebede and Ribka Mekuria

Ethiopian Institute of Agricultural Research, Wondo Genet Agricultural Research Center, Shashemane, Ethiopia

*Corresponding author: Melkamu Hordofa Sigaye, Ethiopian Institute of Agricultural Research, Wondo Genet Agricultural Research Center, Shashemane, Ethiopia, E-mail: hordofa.mel@gmail.com

Received date: March 1, 2022, Manuscript No. AJPSKY-22-12600; Editor Assigned date: March 3, 2022, PreQC No. AJPSKY-22-12600(PQ); Reviewed date: March 17, 2022, QC No. AJPSKY-22-12600; Revised date: March 23, 2022, Manuscript No. AJPSKY-22-12600 (R); Published date: April 01, 2022, DOI: 10.36648/2249-7412.12.4.218

Citation: Sigaye MH, Kebede K, Mekuria R (2022). Response of Tef [Eragrostis tef (Zuccagni) Trotter] to Nitrogen and NPS-B Blended Fertilizers on Andisols of Southern Ethiopia. Asian J Plant Sci Res Vol.12 No.4: 218

ABSTRACT

Inorganic fertilizers have been the important tools to overcome soil fertility problems and also responsible for a large part of the food production increases. The study was aimed to determine the optimum level of NPS-B and by supplementing N from urea rates for maximum yield of tef production and to determine economically optimum level of NPS-B and by supplementing N from urea fertilizer. The treatments were: (100 kg ha-1 NPSB+100 kg ha-1 urea), (150 kg ha-1 NPSB+100 kg ha-1 urea), (200 kg ha-1 NPSB+100 kg ha-1 urea), (150 kg ha-1 NPSB+200 kg ha-1 Urea), (150 kg ha-1 NPSB+200 kg ha-1 urea), (150 kg ha-1 NPSB+200 kg ha-1 urea), (200 kg ha-1 NPSB+200 kg ha-1 urea), (250 kg ha-1 NPBS+200 kg ha-1 urea), (100 kg ha-1 NPSB+300 kg ha-1 Urea), (150 kg ha-1 NPSB+300 kg ha-1 urea), (200 kg ha-1 NPSB+300 kg ha-1 urea), (250 kg ha-1 NPSB+300 kg ha-1 urea), control and R-NP (60N 35 P2O5). The treatments were arranged in a randomized complete block design and replicated three times. The maximum biomass yield (13.2 tone ha-1), grain yield (2.6 tone ha-1) was obtained from the application of 200 kg ha-1 of NPS-B blended fertilizer with 100 kg ha-1 urea. The economic analysis revealed that the highest net benefit of 93885.9 ETB ha-1 with marginal rate of return (MRR%) of 143.8% was obtained in response to the application of 200 kg ha-1 of NPS-B blended with 100 kg ha-1 of urea. However, the lowest net benefit was obtained from an unfertilized or control plot. Therefore, applications of 200 kg ha-1 NPSB of blended plus 100 kg ha-1 of urea is economically advisable for farmers in the Halaba districts southern, Ethiopia and areas with similar agro-ecological and soil types for better tef production.

Keywords: Weed; Wheat; Herbicide; Evaluation; Hadiya

Introduction

Teff (Eragrostis tef /Zucc. / Trotter) is a versatile cereal crop concerning adaptation for the diverse agro-climatic and soil conditions. For this reason, about 2.8 million hectares of land are cultivated annually, covering more than 28% of the total area annually under cereals in Ethiopia. However, its grain yield productivity level is generally low and equal to 1.15 Mg ha–1. Nitrogen and phosphorus nutrients are the major tef yield-limiting factors, the unbalanced and un-optimal fertilization of Ethiopian soils by applications of only DAP and urea (N and P containing fertilizers) for a long period has led to severe nutrient mining of the agricultural soils, particularly when the entire crop biomass (grain and stover) is removed from the land. Such low levels of tef yield are widely believed to be due to low soil fertility caused by low and unbalanced fertilizer application [1].

On the other hand, continuous application of DAP (Di-Ammonium Phosphate: 18%-46% N-P2O5) containing only Nitrogen (N) and Phosphorus (P) without due consideration of other nutrients is known to cause depletion of secondary and micronutrients. Recently, the agricultural extension program has promoted a blanket recommendation of 100 kg DAP and 100 kg urea/ha for all cereal crops and soil types but the actual application rate is 65 kg DAP and 45 kg urea/ha. The low and unbalanced fertilizers application together with poor soil fertility management is presented as the major cause for low agricultural productivity in Ethiopia [1]. Under such conditions, the application of multinutrient blended fertilizers is acknowledged for being able to enhance the productivity and nutrient use efficiency of crops.

Recently, according to the soil fertility map Ethiopia soil analysis data revealed that the deficiencies of most of the nutrients such as nitrogen (86%), phosphorus (99%), sulfur (92%), born (65%), zinc (53%), potassium (7%), copper, manganese, and iron were widespread in Ethiopian soils. Similarly, Asgelil et al. found that the soil analyses and site-specific studies also indicated that elements such as K, S, Ca, Mg and micronutrients (Cu, Mn, B, Mo and Zn) were becoming depleted and deficiency symptoms were observed in major crops in different parts of the country. Inorganic fertilizers have been an important tool to overcome soil fertility problems and are also responsible for a large part of the food production increases.

The drive for higher agricultural production without balanced use of fertilizers created problems of soil fertility exhaustion and plant nutrient imbalances not only of major but also of secondary macronutrient and micronutrients. Similarly [3,4] stated that the deficiencies of secondary macronutrients and micronutrients will arise if they are not replenished timely under intensive agriculture. Consequently, to overcome this problem, multi-nutrient balanced fertilizers containing N, P, K, S, B and Zn in blended form have been issued to ameliorate site-specific nutrient deficiencies and thereby increase crop production and productivity.

Having considered the problems outlined above, the Ethiopian government has been promoting the use of multinutrient blend fertilizers since 2015. The promotion of blend fertilizer follows from the results of the soil fertility survey and preparation of the regional nutrient deficiency atlas of the country under the Ethiopian Soil Information System project. To supply sulfur and Boron commercial fertilizer, DAP is replaced by NPSB. Since the composition of newly introduced fertilizer differs from that of familiar fertilizer (DAP), the appropriate rate is not determined, and insufficient information for Tef production in the study area. Therefore, this research aimed to determine the optimum level of NPS-B by supplementing N from urea for maximum yield of tef production and to determine the economically optimum level of NPS-B by supplementing N from urea fertilizer at Halaba district, Southern Ethiopia.

Methods

Description of the experimental area

The study was conducted during the 2019-2020 cropping season in the Halaba districts of Southern Ethiopia. The geographic locations of the experimental sites in Halaba (7°24'53.87" N and 38°6'55.54" E at an altitude of 1790 m.a.s.l.). The major crops are grown in the area where maize, wheat, sorghum, barley, tef, and pepper. The dominant soil type in the study area in Andisols.

Experimental set-up and procedure

The experimental sites were prepared for sowing using standard cultivation practices and were plowed using oxendrawn implements. The experiment was laid out in randomized complete block design with three replicates for each treatment and detail of the treatments (Table 1). The blended and phosphorus-containing fertilizers from NPSB and Triple Superphosphate (TSP) respectively were basally applied once at sowing to minimize losses and increase use efficiency. Tef (Quncho DZ-Cr-387 variety) was used. The nitrogen fertilizer from Urea was applied in the row two times; half at sowing and the other half during the maximum growth; between the third and fourth weeks of sowing, and at the full tillering stage. All other agronomic management practices were applied as per recommendation. The necessary data were collected at the right time and crop growth stage.

Crop, soil data sampling and analysis

The crop yield and yield components collected included several effective tillers, days to 50% flowering, days to panicle emergence, days to maturity, plant height, panicle length, grain yield, biomass yield and seed weight. Representative composite surface soil samples were collected from 0 cm-20 cm depth at each experimental unit just before sowing. After manual homogenization, the samples were ground to pass a 2 mm sieve. Soil particle size distribution was determined by the Boycouos hydrometric method [5]; pH of the soils was measured in water suspension

in a 1:2.5 (soil: water ratio); organic carbon was determined using the wet oxidation method [6]; total nitrogen was determined using Kjeldahl digestion with concentrated H2SO4 and K2SO4-catalyst mixture [7] available P was determined using the Olsen method [8]; total sulfur in soil extracts was done using Turbidimetric method. The cation exchange capacity was determined after extracting the soil samples by ammonium acetate method (1N NH4O Ac) at pH 7.0 Exchangeable Acidity (EA) Al+3 and H+ were determined from a neutral 1N KCl extracted solution through titration with a standard NaOH solution.

Table 1: Detail of treatment set up and nutrient levels. The nutrients level of in 100 kg of NPS-B were (18.1 N-36.1 P2O5 -0.0 K2O+6.7 S+0.0 Zn+0.71 B).

No	Urea (kg ha 1)	NPS-B (kg ha 1)	Nutrient level (kg ha 1)				
			Ν	P2O5	S	В	
1	Control	0	0	0	0	25	
2	Rec-N and P	60	35	0	0	25	
3	100	100	64.1	36.1	6.7	0.71	
4	100	150	73.15	54.15	10.05	1.065	
5	100	200	82.2	72.2	13.4	1.42	
6	100	250	91.25	90.25	16.75	1.775	
7	200	100	110.1	36.1	6.7	0.71	
8	200	150	119.15	54.15	10.05	1.065	
9	200	200	128.2	72.2	13.4	1.42	
10	200	250	137.25	90.25	16.75	1.775	
11	300	100	156.1	36.1	6.7	0.71	
12	300	150	165.15	54.15	10.05	1.065	
13	300	200	174.2	72.2	13.4	1.42	
14	300	250	183.25	90.25	16.75	1.775	

Economic analysis

Economic analysis was performed to investigate the economic feasibility of the treatments. Partial budget and marginal analyses were used. Current prices of tef, urea, TSP and NPS-B fertilizer were used for the analysis. The potential response of crops towards the added fertilizer and price of fertilizers during planting ultimately determine the economic feasibility of fertilizer application [9]. The market cost of tef was 42.00 Eth-birr kg-1. The prices for blended fertilizers NPSB, TSP and Urea were 20.50, 20.50 and 18.25 Eth-birr kg-1 respectively. The cost of other production practices like seed and weeding were assumed to remain the same or insignificant among the treatments.

Analysis of the Marginal Rate of Return (MRR) was carried out for non-dominated treatments, and the MRRs were compared to a Minimum Acceptable Rate of Return (MARR) of 100% to select the optimum treatment [9]. The net benefit per hectare for each treatment is the difference between the gross benefit and the total variable costs. The average yield was adjusted downward by 10% to reflect the difference between the experimental field and the expected yield at farmers' fields and with farmer's practices from the same treatments.

Statistical analysis

Data from the field and laboratory were tested for normality, before being subjected to Analysis of Variance (ANOVA) using SAS software program version 9.4. The significant difference among treatment means was evaluated using the least significant difference at ($p \le 0.05$).

Results

Physicochemical properties of the soil

The soil particle size distributions of the experimental site were 63.1%, 21.3% and 15.3% sand, silt and clay respectively. Thus, the soil textural class of the soil was sandy loam. The pH value of the soil was 6.5 (1:2.5 soil: water). The soil pH has a vital role in determining several chemical reactions in influencing plant growth by affecting the activity of soil microorganisms and altering the solubility and availability of most of the essential plant nutrients, particularly the micronutrients such as Fe, Zn, B, Cu and Mn [10]. (Sumner, 2000). The available P content of the soil was 11.2 (ppm). According to Olsen et al. [9]. rating, the status of available P of both sites categorized under medium level (Table 2). The organic carbon and total nitrogen content of the experimental soil were (1.5%) and (0.12%) respectively. According to Tekalign [11]. Rating of organic carbon and total nitrogen content of the experimental site soil under moderate to medium. Total sulfur (S) (2.8) and statuses of the soil found were d under the low category [12] (Table 2). Whereas the cation exchange capacity of soil sites, 12.8 meq 100g sample-1, respectively, London, [13] Hazelton and Murphy [14] were found at medium status (Table 2).

Table 2: Physio-chemical properties of the soil.	Effects of NPS-B b	blended fertilizers an	nd urea on yield an	nd yield
components of tef				

Parameters	Value	Rating	Reference
Sand (%)	63.1		-
Silt (%)	21.6		
Clay (%)	15.3		
Textural class	Sandy Loam		[16]
pH(1:2.5 soil: water)	6.1		
Ava P (mg kg-1)	11.2	Medium	[11]
S (mg kg-1)	2.8	Low	[19]
TN (%)	0.12	Low	[18]
OC (%)	1.5	Low	[18]
CEC (cmol (+) kg-1)	12.8	Moderate	[20,21]

Application of NPSB blended fertilizers and Urea were highly significantly (p<0.01) influence yield and yield components of tef. The pooled mean analysis revealed that the tallest plant height (100.9 cm) was obtained from the application of 250 kg ha-1 NPSB with 300 kg ha-1 urea. The longest spike length (36.9 cm) was obtained from the application of 200 kg ha-1 NPSB with 300 kg ha-1 urea. And the maximum number of tillers per m2 (385) was obtained from the application of 200 kg ha-1 NPSB with 100 kg ha-1 urea. However, the minimum tef yield components were obtained from unfertilized or control plots.

This significant increment of tef yield components may be attributed to the fact that N usually favors vegetative growth of teff, resulting in higher stature of plants while P is the main element involved in energy transfer for cellular metabolism in addition to its structural role and Sulphur enhanced the formation of chlorophyll and encouraged vegetative growth [15]. Similarly, an adequate supply of phosphorus increases tiller emergence, especially secondary tillers and their survival, it helps in increasing the biomass yield through proper regulation of carbohydrates translocation. An ample supply of boron facilitates photosynthetic activities and leaf expansion that leads to improved plant growth [16]. This might be due to plants grown on plots treated with a higher rate of N for their vegetative growth, higher P phosphorus for their good root development, higher level of S for a high number of tillers and B for its higher cell division; it also contributed to increasing the total number of tillers per plant and influenced the straw yield [17]. The current findings were in line with [18] were reported that the contained application of blended fertilizer with urea provides significantly higher yield components on tef.

The combined year's analysis revealed that application of NPSB blended fertilizer with N from urea was highly (p<0.01) significantly influence the biomass and grain yield (Tables 3-5). The maximum biomass yield (12.9 tone ha-1), grain yield (2.5 tone ha-1) was obtained from the application of 200 kg ha-1 NPSB blended fertilizer with 100 kg ha-1 urea (Figure 5). The total grain yields obtained at 200 kg ha-1 NPSB blended with 100 kg ha-1 urea rates of fertilizers exceeded yield produced by 137.1% and 70.2%, unfertilized or control and recommended NP fertilizer respectively.

Tef yields were consistently increased across blended NPS-B fertilizer and N from urea. This might be due to better crop growth rate, leaf area index, and accumulation of photo-assimilate due to maximum days to maturity by the crop, which ultimately produced more biomass and grain yields. Fageria et al. [17] also indicated that the application of S enhanced the photosynthetic assimilation of N in crops. Hence, the application of N and S increased the net photosynthetic rate which in turn increased the dry matter as 90% of dry weight is considered to be derived from products formed during photosynthesis. This result in conformity with this finding of different scholars [18-21] were reported that the application of N from urea was increased biomass, grain of tef.

Table 3 : Mean values of yield and yield components of tef as influenced by nitrogen and NPS-B blended fertilizer during 2019 cropping season. Means with the same letter along the column are not significantly different at $p \le 0.05$, where; PH: Plant height; SL: Spike length; NT: number of tillers; BM: Biomass; GY: Grain yield; R-NP: Recommended rate of Nitrogen and Phosphorous

Treatments (kg ha-1)		DII (am)		NT (2)	DM (4 h = 1)	OV(1 + 1)
Urea	NPSB	PH (cm)	SL (cm)	NI (m2)	BNI (t na-1)	GY (t na-1)
Control		62.1 ^b	13.7 ^b	75°	6.2.0°	0.6 ^h
RNP		99.1ª	33.5ª	354 ^{ab}	13.1 ^{ab}	2.2 ^b
100	100	90.0a	32.4ª	219 ^d	9.4 ^d	1.3 ^{gh}
100	150	95.7ª	33.9ª	277 ^{bcd}	11.7 ^{bc}	2.0 ^{cdd}
100	200	100.8ª	32.1ª	385ª	13.5ª	2.8ª
100	250	100.3ª	34.3ª	324 ^{abc}	12.4 ^{ab}	2.0 ^{bc}
200	100	93.5ª	35.3ª	318 ^{abc}	10.8 ^{bcd}	1.9 ^{bcd}
200	150	100.0ª	34.2ª	275 ^{bcd}	12.2 ^{abc}	1.9 ^{bcd}
200	200	92.0ª	28.5ª	241 ^{cd}	9.8 ^{cd}	1.7 ^{cde}
200	250	100.0ª	34.5ª	248 ^{cd}	12.7 ^{ab}	2.0 ^{bc}
300	100	99.1ª	33.1ª	343 ^{ab}	12.3 ^{abc}	1.8 ^{bc}
300	150	97.8ª	32.2ª	315 ^{abc}	12.6 ^{ab}	1.6 ^{cde}
300	200	103.0ª	36.9ª	308 ^{abcd}	13.0 ^{ab}	2.2
300	250	100.3ª	33.0ª	296 ^{abcd}	12.1 ^{abcd}	11 ^g
CV		9.2	21.4	19.1	13.1	9.1
LSD@ <0.05		14.7*	14.7**	91.2**	2692.2**	271.6**

Table 4 : Mean values of yield and yield components of tef as influenced by nitrogen and NPS-B blended fertilizer during 2020 cropping season. Means with the same letter along the column are not significantly different at p ≤ 0.05 , where; PH: Plant height; SL: Spike length; NT: Number of tillers; BM: Biomass; GY: Grain yield, R-NP: Recommended rate of Nitrogen and Phosphorous

Treatments (kg ha-1)		DIL (and)	CL (and)	NT (2)	DM (4 h = 1)	$CV(4 h \cdot 1)$	
Urea	NPSB	PH (cm)	SL (cm)	NI (m2)	BM (t na-1)	GY (Ulla-1)	
Co	ntrol	54.9 ^f	14.9°	99°	5.6 ^d	0.3 ^f	
R	NP	95.2 ^{abc}	25.0 ^{abc}	256 ^{abc}	11.5 ^{abc}	1.7 ^{bcd}	
100	100	82.8°	16.2 ^{de}	154 ^{ef}	9.0 ^{bc}	1.1°	
100	150	88.5 ^{bcde}	24.4 ^{abc}	279 ^{ab}	11.4 ^{abc}	1.7 ^{bcd}	
100	200	96.9 ^{ab}	29.2ª	321ª	12.9ª	2.5ª	
100	250	96.4 ^{ab}	24.8 ^{abc}	226 ^{bcde}	12.8ª	1.9 ^b	
200	100	86.3 ^{cde}	25.8abc	220 ^{bcde}	10.2 ^{abc}	1.6 ^{bcd}	
200	150	96.1 ^{ab}	24.7 ^{abc}	244.0 ^{bcd}	11.9 ^{ab}	1.6 ^{bcd}	
200	200	84.8 ^{de}	20.7 ^{cd}	177 ^{de}	8.6 ^{cd}	1.5 ^{cde}	
200	250	92.8 ^{bcd}	25.0 ^{abc}	250 ^{abcd}	12.7ª	1.7 ^{bcd}	
300	100	91.9	23.6 ^{bc}	245 ^{bcd}	11.7 ^{abc}	1.5 ^{cde}	
300	150	93.9 ^{abcd}	22.7 ^{bc}	217 ^{bcde}	12.1 ^{ab}	1.3 ^{de}	
300	200	95.8 ^{ab}	27.4 ^{ab}	210 ^{bcde}	12.5ª	1.7 ^{bc}	
300	250	103.1ª	23.5 ^{bc}	198 ^{cde}	11.5 ^{abc}	0.6 ^f	
CV		6.3	13.9	20.1	17.1	16.6	
LSD@ ≤ 0.05		9.4*	5.9**	74.5**	3168.3**	412.9**	

Table 5: Pooled mean values of yield and yield components of tef as influenced by nitrogen and NPS-B blended fertilizer during 2019-2020 cropping season. Means with the same letter along the column are not significantly different at $p \le 0.05$, where; PH: Plant height; SL: Spike length; NT: Number of tillers; BM: Biomass; GY: Grain yield: R-NP: Recommended rate of Nitrogen and Phosphorous

Treatments (kg ha-1)		DII (am)	SL (am)	NT (m2)	DM (the 1)	CV(tho 1)
Urea	NPSB	rn (cm)	SL (cm)	IN I (III2)	DM (t na-1)	GY (t lla-1)
Control		58.5°	14.3 ^d	87 ^f	5.9 ^d	0.5 ⁱ
R-NP		97.1 ^{abc}	29.3 ^{abc}	305 ^{ab}	12.3 ^{ab}	1.91 ^{bc}
100	100	86.4°	24.3°	187°	9.3°	1.2 ^g
100	150	92.1 ^{bcde}	29.1 ^{abc}	278 ^{bc}	11.5 ^{ab}	1.8 ^{bcde}
100	200	98.8 ^{ab}	30.7 ^{ab}	353ª	13.2ª	2.6ª
100	250	98.3 ^{abc}	29.6 ^{abc}	275 ^{bc}	12.6ª	2.0 ^b
200	100	89.9°	30.5 ^{abc}	269 ^{bc}	10.5 ^{bc}	1.8 ^{bcde}
200	150	98.0 ^{abc}	29.5 ^{abc}	259 ^{bcd}	12.1 ^{ab}	1.7 ^{cde}
200	200	88.4 ^{cde}	24.6 ^{bc}	209 ^{de}	9.2°	1.6 ^{ef}
200	250	96.4 ^{abcd}	29.8 ^{abc}	249 ^{bcd}	12.7ª	1.8 ^{bcde}
300	100	95.5 ^{bcde}	28.4 ^{abc}	294 ^{bc}	12.1 ^{ab}	1.7 ^{def}
300	150	95.9 ^{abcd}	27.5 ^{abc}	266 ^{bc}	12.3 ^{ab}	1.5 ^f
300	200	99.4 ^{ab}	32.2ª	259 ^{bcd}	12.7ª	1.9 ^{bcde}
300	250	101.7ª	28.3 ^{abc}	247 ^{cd}	11.9 ^{ab}	0.8 ^h
CV		8.1	19.7	19.3	15.3	12.6
LSD@<0.05		8.6*	6.3*	56.7*	1999.2**	237.5**

Economic analysis

As indicated in Table 6, the highest net benefit of 93885.9 ETB ha-1 with a marginal rate of return (MRR%) of 143.8% was obtained in response to the application of 200 kg ha-1 of blended NPSB with 100 kg ha-1 of urea. However, the lowest net benefit was obtained from the unfertilized or control plot (Table 6). Thus, applications of 200 kg ha-1 NPSB of blended plus 100 kg ha-1 of urea is economically advisable for farmers in the study area for better tef production; beneficial as compared to the other treatments in the study area because the highest net benefit and the marginal rate of return were above the minimum level (100%).

Table 6: Nitrogen and NPS-B blended fertilizer effects on partial budget and marginal rate of return analysis for tef production. R-NP: Recommended rate of Nitrogen and Phosphorous, GY: Grain yield; TCV: Total Cost that Varies; NBC: Net Benefit-Cost, GB: Growth Benefit, Marginal rate of return; d: Dominance.

Treatments	GY (t ha-1)	GB	TVC	NBC	MRR%
1	0.5	17694.2	0	17694.2	
2 (R-NP)	1.91	73184.6	3643.5	69541.1	14.2
3	1.2	45552.8	3875	41677.8	d
4	1.8	68350	4900	63450	21.2
7	1.8	67223.5	5700	61523.5	d
5	2.6	99810.9	5925	93885.9	143.8
8	1.7	65435.6	6725	58710.6	d
6	1.8	74862.9	6950	67912.9	40.9
11	1.7	63383	7525	55858	d
9	1.4	60778.6	7750	53028.6	d
12	1.5	56352.2	8550	47802.2	d
10	1.8	69245.8	8775	60470.8	56.3
13	1.9	70213.5	9575	60638.5	0.2
14	0.8	29899.8	10600	19299.8	d

Conclusion and Recommendation

C practice for increasing tef yield component and yield. The result of the economic analysis showed that the combined application of 200 kg ha-1 of NPSB and 100 kg ha-1 of urea gave economic benefit. Therefore, it could be concluded that the application of 200 kg ha-1 of NPSB with supplement 100 kg ha-1 of urea fertilizer combinations was producing economically profitable tef production in the area.

Acknowledgments

We acknowledge the support of the Ethiopian institute of agricultural research and Wondo genet agricultural research center during part of this work.

Conflict of Interest

The authors declare that they have no conflict of interest

REFERENCES

- 1. Elias E, Okoth PF, Smaling EMA (2019) Explaining bread wheat (Triticum aestivum) yield differences by soil properties and fertilizer rates in the highlands of Ethiopia. Geoderma 33(9) (2019) 126–133.
- 2. Hailu H, Mamo T, Keskinen R, Karltun E, Gebrekidan H, et al. (2015) Soil fertility status and wheat nutrient content in Vertisol cropping systems of central highlands of Ethiopia. Agric Food Secur 4: 19.
- 3. Fageria NK, Baligar VC (2001) Improving nutrient use efficiency of annual crops in Brazilian acid soils for sustainable crop production. Commun Soil Sci Plant Anal 32: 1303-1319.
- 4. Singh J, Bisen S, Bora DK, Kumar R, Bera B (2011) Comparative study of organic, inorganic and integrated plant nutrient supply on the yield of Darjeeling tea and soil health. Field Crop Research 58: 58-61. [Crossref], [Google Scholar]
- Van Reeuwijk LP (2002) Procedures for Soil Analysis FAO, International soil reference and information center, 6700 AJ Wageningen, The Netherlands. (6th Ed), pp: 120.
- 6. Walkley A, Black I (1934) An examination of different methods for determining soil organic matter and a proposed modification of the chromic acid titration method. Soil Science 37: 29-37
- 7. Black CA (1965) Methods of soil analysis. Part 2, Am Soc Agronomy. Madison, Wisconsin, USA, pp: 56-60.
- 8. Olsen SR, Sommers LE (1982) Phosphorus. Methods of soil analysis, Amer Soc Agronomy, Madison, Wisconsin. Part 2, (2nd Ed), pp: 403-429. [Crossref], [Google Scholar]
- 9. CIMMYT (1988) From agronomic data to farmer's recommendations: An economics training manual. CIM-MYT, Mexico, DF, pp: 79.
- 10. Sumner ME (2000) Handbook of soil science., CRC Press, Boca Raton, FL, pp: 88.
- 11. Tadese T (1991) Soil, plant, water, fertilizer, animal manure, and compost analysis. Working document international livestock research center for Africa, Addis Ababa, Ethiopia, 13.
- 12. Ackerman D, Large BSR (2015) The agronomy handbook. Midwest laboratories, pp34.
- 13. London JR (1991) Booker tropical soil manual hand-book for soil survey and agricultural land evaluation in the tropical and subtropics. Long man scientific and technical, ESSXEX, New York. pp: 474.
- 14. Hazelton P, Murphy B (2007) Interpreting soil test results: What do all the numbers mean? 2nd Edition. CSIRO Publishing. pp: 152.
- 15. Havlin JL, Beaton JD, Tisdale SL, Nelson WL (2005) Soil fertility and fertilizers: An introduction to nutrient management. Pearson Prentice Hall, Upper Saddle River, New Jersey.
- 16. Tahir M, Tanveer A, Shah T, Fiaz N, Wasaya A (2009) Yield response of wheat (Triticum aestivum l.) to Boron application at different growth stages. Pak J Life Soc Sci 7: 39-42. [Crossref], [Google Scholar], [Indexed]
- 17. Fageria NK, Baligar VC, Jones CA (2011) Growth and mineral nutrition of field crops. 3rd Edn, Taylor & Francis Group, New York, 52: 530.
- Wakjira T (2018) Teff yield response to NPS fertilizer and methods of sowing in East Shewa, Ethiopia. J Agric Sci 13: 162-173.
- 19. Gebretsadik H, Haile M, Yamoah, CF (2009) Tillage frequency, soil compaction, and N-fertilizer rate effects on yield of tef [Eragrostis tef (Zucc.) Trotter]. Ethiop J Sci 1: 82-94. [Crossref], [Google Scholar], [Indexed]

- 20. Haileselassie B, Habte D, Haileselassie M, Gebremeskel G (2014) Effects of mineral nitrogen and phosphorus fertilizer on yield and nutrient utilization of bread wheat (Tritcum aestivum L.) on the sandy soils of Hawzen District, Northern Ethiopia. Int J Agr For Fish 3: 189- 198. [Crossref], [Google Scholar], [Indexed]
- 21. Tekle L, Wassie H (2018) Response of tef (Eragrostis tef (Zucc.) Trotter) to blended fertilizers in Tembaro, Southern Ethiopia. J Biol Agri Health care 8: 34-39. [Crossref], [Google Scholar]