

Risk Factors of Osteoarthritis in Women

Salah Alghanem*

Department of Gynaecology, Beth Israel Deaconess Medical Centre, Harvard Medical School, Boston, Massachusetts, USA

*Corresponding author: Salah Alghanem, Department of Gynaecology, Beth Israel Deaconess Medical Centre, Harvard Medical School, Boston, Massachusetts, USA, E-mail: salah@healthcore.com

Received date: September 3, 2021; Accepted date: September 17, 2021; Published date: September 24, 2021

Citation: Alghanem S (2021). Risk Factors of Osteoarthritis in Women. J Women's Health Reprod Med Vol.5 No.5:e021.

Description

Osteoarthritis is the most common form of arthritis, which affects millions of people around the world. Symptomatic knee Osteoarthritis occurs in 10% men and 13% in women aged 60 years or older. It happens when the protective cartilage commonly the ends of the bones are worn over time. Although the osteoarthritis can damage any joint, the disorder most commonly affects the joints in their hands, knees, hips and spine. The symptoms of osteoarthritis can be generally administered, although the damage to the joints can be reversed. Staying active, keeping a healthy weight and receiving certain treatments could slow down the progression of the disease and help improve the pain and function of the osteoarthritis is the most common chronic articulation condition (long term).

A joint is where two bones come together. The ends of these bones are covered by protective fabric called cartilage. With osteoarthritis, this cartilage breaks, which makes the bones together inside the joint break. This can cause pain, rigidity and other symptoms occur more frequently, although it can occur in adults of any age. Osteoarthritis is also called degenerative articulation disease, degenerative arthritis and arthritis to wear. Here, all women need to know about osteoarthritis, from treatment to prevention and more. Cartilage is a hard and rubbery substance which is flexible and softer than the bone. Women task is to protect the ends of the bones inside a joint and allow them to move easily with each other. When the

cartilage is divided, these bone surfaces are pungent and harsh. This can cause pain within articulation and irritation in the surrounding tissues. Damaged cartilage cannot be repaired. This is because cartilage does not contain blood vessels.

When the cartilage is completely wearing, the damping buffer supplies disappears, which allows bone contact. This can cause intense pain and other symptoms associated with Osteoarthritis. Here, what else needs to learn about cartilage, joints and osteoarthritis. Osteoarthritis is a progressive condition with five phases, from 0 to 4. The first phase (0) represents a normal joint. Phase 4 represents one or serious. Not all those who have or progress to phase 4. The condition is often stabilized long before reaching this phase. People with serious or have extended or complete loss of cartilage in one or more joints. The bone friction associated with this can cause serious symptoms, such as increased swelling and inflammation. The amount of synovial liquid within the joint can increase. Normally this liquid helps reduce friction during movement. However, in greater quantities, it can cause joint swelling. The fragments of broken cartilage can also float inside the synovial fluid, which increases pain and swelling. Decrease in the range of movement. It is possible that it cannot move also due to rigidity or pain in its joints. This can make you more difficult to enjoy day today activities that come easily. Joints may be less stable. The articular damage caused by or serious or reversible, but treatment can help reduce symptoms. Discover everything you need to know about advanced osteoarthritis.