



SEEING DOUBLE: A CASE OF DOUBLE PRIMARY CANCER: NEUROENDOCRINE TUMOR OF THE OVARY AND ENDOMETRIAL ADENOCARCINOMA IN A 45 YEAR OLD NULLIGRAVID”

Michelle Hubilla

Department of Obstetrics and Gynecology, Makati Medical Center, Makati City, Phillipines

ABSTRACT

A 45 year old nulligravid presented with changes in sensorium after losing balance and hitting her head, with noted symptoms of restlessness, disorientedness and intermittent febrile episodes. Upon work-up, serum calcium and parathyroid hormone were noted to be significantly elevated. Investigating the pathology of hypercalcemia led to an incidental finding of a pelvoabdominal mass. Computed tomography (CT) scan of the whole abdomen showed a heterogeneously enhancing mass noted in the bilateral hemipelvis measuring 8.5 x 5.0 x 6.4 cm in the right and 3.4 x 2.5 x 4.1 cm in the left and pelvic and retroperitoneal lymphadenopathy, for which a neoplastic process was considered, probably gynecologic in origin. With the aforementioned signs and symptoms, patient was managed as a case of Pelvoabdominal mass, to consider Anti-NMDAR encephalitis vs. Parathyroid hormone-secreting tumor vs. Germ Cell tumor of the ovary. Patient underwent extrafascial abdominal hysterectomy with bilateral salpingoophorectomy, bilateral lymph node dissection with frozen section. Intraoperatively, tumor was found to be on the paratubal area of the right fallopian tube. It was then sent for frozen section revealing a histopathologic diagnosis of metastatic high grade papillary adenocarcinoma. Other specimens that were submitted include the cervix, uterus, ovaries left fallopian tube had a final histopathologic diagnosis of a mixed endometrioid and undifferentiated carcinoma with carcinoma involving more than half of the myometrial depth with invasion into the cervix, right and left parametria, right and left fallopian tube and ovaries. Lymph node dissection also revealed an extensive lymphovascular invasion. The specimens were stained with synaptophysin and chromogranin A, two of the most common used neuroendocrine markers. It was then confirmed that there are two separate malignancies present in the patient, the Endometrial Endometrioid Adenocarcinoma (Figo Grade 1) well differentiated, confined to the endometrium and the other, a Large cell Neuroendocrine carcinoma involving the right and left ovaries with extension to bilateral fallopian tube, bilateral parametria and invasion into the myometrium and cervix Metastatic large cell neuroendocrine carcinoma to the pelvic lymph nodes. Neuroendocrine tumors are spectrum of malignancies that arise from the diffuse neuroendocrine cell system that can cause debilitating symptoms and potentially life threatening conditions for patients (Gardner, Reidy-Lagunes & Gehrig, 2011)



Keywords–

Lymphovascular, Endometrioid, Adenocarcinoma, Neuroendocrine

Biography – Michelle O. Hubilla M.D. is a 4th year resident in the Department of Obstetrics and Gynecology in Makati Medical Center. At the age of 23 years old, she completed her degree in Medicine at the De La Salle Health Sciences Institute and her one year internship at the Department of Health hospitals. Currently, she is on her last year of training in one of the top hospitals in the country, The Makati Medical Center, and aspires to be a gynecologic oncologist and serve the Filipino people

[2nd International Conference on Womens Health, Gynecology and Obstetrics, July 06-07, 2020 Webinar](#)

Abstract Citation : [MICHELLE O. HUBILLA, SEEING DOUBLE: A CASE OF DOUBLE PRIMARY CANCER: NEUROENDOCRINE TUMOR OF THE OVARY AND ENDOMETRIAL ADENOCARCINOMA IN A 45 YEAR OLD NULLIGRAVID, WOMENS HEALTH 2020, July 06-07, 2020 Webinar](#)