

## Vertebral fractures among patients referred for Bone densitometry screening in Dubai Primary Healthcare Facilities


**Anood Jamal Alshaali**

Elderly Care Unit, Dubai Health Authority, United Arab Emirates



### Abstract

Vertebral fractures are one of the most common fractures associated with low bone mineral density. However two third to three fourth of patients with vertebral fractures are not clinically recognized. The objective of this study was to determine the prevalence of vertebral fractures in patients referred for bone densitometry and the most common site of fracture. The study was carried out in the osteoporosis clinic in Dubai Primary healthcare centre. A total of 120 patients were examined using the dual energy x-ray absorptiometry. Of all the patients, 48.3% were osteoporotic and 40.9% were osteopenic. The overall prevalence of vertebral fracture was 14.2%. The result showed that the prevalence of vertebral fracture was higher in female compared to male (15.7% and 9.7% respectively). It was found that patients aged 80 and above had the highest prevalence of vertebral fracture (54.5%). Undiagnosed vertebral fractures were common. Therefore, it is crucial to prevent vertebral fracture through early diagnosis and appropriate treatment of osteoporosis



### Biography

Dr Anood Al Shaali has graduated from the Royal College of Surgeons in Ireland. She has completed the Arab Board in Community Medicine in 2013. She is currently working as a Community Medicine Senior Specialist Registrar in the Elderly Care Unit, Dubai Health Authority.

[14th International Conference on Rheumatology and Arthroplasty](#) | London, UK | September 21-22, 2020

**Citation:** Ahmed Ismail, Effect of Insulin Local Injections in Diabetic Patients with Carpal Tunnel Syndrome In Comparison With Corticosteroid Local Injection, Rheumatology 2020, 14th International Conference on Rheumatology and Arthroplasty, London, September 21-22, 2020