

DNA Fingerprinting and CRISPR cas9 System

Umair Masood

Abstract

Leicester university geneticist Alec Jeffrey's develops a technique called DNA finger printing in 1985 it allows the DNA sample from different people to be compared look for similarities and differences. It used the solving crime and can confirm if the people are related to each other like paternity testing. There is section of chromosomes where an instead of gene consisting of a long sequence of bases, they are usually (15-100) base pairs long that are repeated in many times these repeated sequences called Variable Number of Tandem Repeat. we can know the VNTR sequence of any person than we can design a gRNA. Let's say you have a DNA sample with fluorescent labeled from victim and you want to make sure that VNTR sequence you are interested is in match with victim. We can design a CRISPR to scan through DNA or find specific VNTR. The CRISPR scan the DNA if the CRISPR does not find targeted VNTR it does not bind to it its means that no fluorescence color appears under UV-light but its scan and find its target and this binding create a fluorescence signal its means that VNTR can be occur in a DNA.

Keywords: DNA finger printing; CRISPR cas9; UV-light; VNTR sequence; Fluorescence color

Department of Biology, Comsats University, Islamabad

*Corresponding author: Umair Masood

✉ Umairawan0505@gmail.com

Department of Biology, Comsats University, Islamabad

Citation: Umair M (2021) DNA Fingerprinting and CRISPR cas9 System. Eur Exp Biol Vol.11 No. 3:127

DNA Fingerprinting System Parts

CAS9-Protein

Cas9 protein is also called destructive protein Its main function is to cut DNA and thereby alter a cell's genome [1-5].

Guide-RNA (VNTR)

The main part of our technique is gRNA or VNTR the guide RNA is a specific that recognizes the target DNA area of interest and directs the Cas nuclease there for editing (Figure 1).

Template DNA

The template DNA is our target DNA that we want to check either Specific STR is present or not.




Figure 1: Cs9 bind and cut the desire sequence and emit a fluorescence signal.

DNA Fingerprinting and CRISPR Cas9 Protocol

The protocol contains two steps.

1. Labeling of Template DNA
2. CRISPR cas9 working protocol

Labeling of Template DNA

Reagent requires

3. DNA dye 70 μ l
4. Buffer 1:275 μ l
5. Buffer 2:255 μ l
6. AT rich DNA:400 μ l
7. GC rich DNA:400 μ l
8. 50:50 at and GC:500 μ l
9. 100mM NAOH 650 μ l

Procedure

10. Label the tubes 1,2,3 and 4
11. Add 10 μ l buffer to each tube
12. Add 10 μ l dye to each tube
13. Add 5 μ l of DNA sample to tube 1,2 and 3
14. Do not add an any DNA to tube 4 because tube 4 is serve as negative control
15. Mix the reagent by pipetting up and down 3-4 time
16. Enter the heat block parameter such as 95°C for 2 min
17. Run a PCR
18. View under UV-light

CRISPR Cas9 Working Protocol

Procedure

Table 1: a):Position a 30 μ l reaction along micro centrifuge tube on ice with the following sequence; b):Softly mix the reaction mixture and centrifuge it; c):Incubate at 37°C for 20min; d): View under UV-light.

Component	20 μ l Reaction
Template DNA	x μ l (~100 ng)
Guide RNA	x μ l (~4000 ng)
10 X Cas9 Reaction Buffer	3.0 μ l
Cas9 Nuclease	1.0 μ l
water	30.0 μ l

References

1. Mnookin J L, Cole S A, Dror I E, Fisher B A (2010) The need for a research culture in the forensic sciences. *UCLA L Rev* 58:725.
2. Chambers GK, Curtis C, Millar CD, Huynen L, Lambert DM (2014) DNA fingerprinting in zoology: past, present, future. *Invest Gen* 5:1-1.
3. Kimura P, Nakane T, Ishitani R, Hatada I, Zhang F et al. (2014) Molecular mechanism of CRISPR. *J Appl Crystallogr* 156:935-949.
4. Blum B, Simpson L (1990) Guide RNAs in kinetoplastid mitochondria have a nonencoded 3' oligo (U) tail involved in recognition of the preedited region. *Cell* 62:391-397.
5. Amann R, Ludwig W (2000) Ribosomal RNA-targeted nucleic acid probes for studies in microbial ecology. *FEMS Microbio Rev* 24:555-565.