

Successful 30 years follow up after surgery for dystopic lingual goiter in 2 adolescents

Romanchishen Anatoly F., Vabalayte Kristina, Karpatsky Igor V

Saint-Petersburg State Pediatric Medical University, Russia



Abstract

Introduction: During embryonic development thyroid tissue gradually falls down from tongue root to the typical thyroid gland's place. Lingual Goiter emerge in 1 patients of 100.000 population and in 1 (0.03%) of 3000 Thyroid Diseases patient. We observed 8 (0.025%) lingual goiter (LG) of 32022 thyroid diseases patient (1973 - 2019) and 3 of them were operated on. As a result of thyroid tissue (TT) omitting LG, supra – praehyoid - behind and infrahyoid goiter of thyroid remnants along of thyroglossal duct could appeared with (ectopia) or without (dystopia) presence TT in typical place. Indications for Surgical treatment of a LG patients: relapse or profuse bleeding, aggressive grooves of LG with dyspnea and dysphagia; suspicions of malignancy, thyrotoxic LG. Thyroid disorders are one of the most common endocrine problems in children and adolescents. Children with thyroid disorders usually present with an enlargement of the thyroid gland (goitre), with or without symptoms of thyroid hormone deficiency or excess. These symptoms are generally insidious in onset, which may delay diagnosis by several weeks to several months. Because a goitre is often the first sign of thyroid disease, the diagnostic considerations can be approached from the perspective of the goitre. By far, the most frequent cause of a goitre is autoimmune thyroid disease, although a benign colloid goitre is also a common cause. The present review focuses on the diagnostic approach to a child with a hypothyroid, hyperthyroid or euthyroid goitre. The most common finding related to thyroid disease is a diffuse enlargement of the thyroid, with the right lobe being frequently larger than the left lobe. The enlargement of the thyroid is generally mediated by an increase in the pituitary-derived thyroid stimulating hormone (TSH) or in antibodies that bind to the TSH receptor, such as the thyroid stimulating immunoglobulins (TSIs) found in Graves disease. Inflammation or infiltration may cause diffuse, symmetrical enlargement, although the gland is usually asymmetric and nodular. The most common inflammatory process is autoimmune thyroiditis

Keywords: Goitre, Hyperthyroidism, Hypothyroidism, Thyroid nodule

Material and Method: Patient 39 years old woman with lingual, suprahyoid and praehyoid thyroid gland goiter (1997). She had tumor mass in the root of tongue, supra - and praehyoid part of the neck from 10 years old. Since 1994 the tumors has increased in size quickly and caused of dyspnea, dysphagia and

nasonnement voice changing. CT scans of the patient neck has showed: LG narrowed of oropharings, TT is absent in typical place. Surgery of patients (August 08. 1997). submandibular approach has find LG, suprahyoid, praehyoid goiters and was take it out 04. 08. 1997. 2.5 month later after patients operation was performed Tc99m scintigram - radiodrug accumulation was absent on the neck. She takes the replacement LT4 therapy.

Conclusion: 1. Wait-and-see policy of lingual goiter patients treatment can be dangerous because suffocation might occur. 2. Lingual goiter would be successfully removal through neck submandibular approach with lingual root resection.



Biography:

Anatoly F Romanchishen is a Professor of Surgery and Oncology, Chief of Hospital Surgery and Oncology Department of Saint-Petersburg State Pediatric Medical Academy, Director of St. Petersburg Center of Endocrine Surgery and Oncology, Member of American, European, Asian, and Italian (Honorable), Associations of Endocrine Surgeons, Honored Doctor of Russian Federation.

Speaker Publications:

1. "P298 Thyroid cancer patient's fertility after radioactive iodine therapy"; Archives of Disease in Childhood. / 2019 / 104(3) / pp A277.1-A277.
2. "OC54 Intraoperative nerve integrity monitor as part of safety dissections of recurrent laryngeal nerve in child and adolescents thyroid surgery"; Archives of Disease in Childhood / 2019 / 104(Suppl 3) / A22.3-A23.
3. "Prophylactic thyroidectomy as method of medullary thyroid carcinoma prevention in children from MEN syndrom families"; Pediatrician (St. Petersburg) / 2017 / Vol 8(5), pp 5-11.

4. "Russian Thyroid Surgery Origin, Development in XIX - Beginning of XX Century and Present Time"; Endocrinology Surgery/ 2014 / Vol 8(1), pp 27-36.
5. "Bilateral synchronous parathyroids cancer and femoral neck fracture as the complications of tertiary hyperparathyroidism"; Endocrine Surgery / 2014 / Vol 7(4), pp 21-24.

[32nd European Pediatrics Congress](#); Webinar; July 22-23, 2020.

Abstract Citation:

Anatoly Romanchishen, Successful 30 years follow up after surgery for dystopic lingual goiter in 2 adolescents, Euro Pediatrics 2020, 32nd European Pediatrics Congress; Webinar; July 22-23, 2020

(<https://europediatrics.pediatricsconferences.com/abstract/2020/successful-30-years-follow-up-after-surgery-for-dystopic-lingual-goiter-in-2-adolescents>)