

Challenges and opportunities in accessing veritable vaccines, and curbing the dissemination and excesses of the SARS-CoV-2 or COVID-19 variants

Dr Chrysanthus Chukwuma Sr

The Chrysanthus Centre for Future-Oriented Studies, CTR Future-Oriented Ltd Abakaliki, Ebonyi State, Nigeria

Email-cfos_nigeria@yahoo.com, Tel. No.: +2348075011261

Abstract

Viruses including SARS-COV-2 that is the etiologic agent of the COVID-19 pandemic, spatiotemporally evolves or mutates to its variant with potentialities for adverse opportunistic infections via replications and transmissions or disseminations. The dissemination and transmission of the Delta variant is accelerating more than expected partly due to unvaccinated persons not treating the pandemic as seriously as necessary. The test and tracing service must be effected because a vast proportion of the population is unprotected. It is crucial to restrict movements. Vaccinated persons must socialize indoors in well-ventilated ambients, and avoid non-essential foreign travels. The evolution of the COVID-19/SARS-CoV-2 pandemic portends a well-nigh impossible termination of events. It is, therefore, imperative to focus on the cost and not merely the opportunity cost to control the never-ending SARS-CoV-2/COVID-19 pandemic. We need to make the public health system a priority with emphasis on cost sharing and cost saving for vaccines and other healthcare modalities to reach the poor and the vulnerable populations in developing countries and elsewhere for safety and protection of lives through enhanced substantial, restorative and sustainable financing, technical expertise and capacity building.

Keywords: spike protein, mutation, dissemination, transmissibility, COVID-19, SARS-CoV-2, social restrictions

Introduction

As the scourge of the COVID-19 or SARS-CoV-2 [1] pandemic remains unabated, it becomes imperative to configure the public health system as a priority with emphasis on cost saving and cost sharing for vaccine distribution and consumption to extend to the poor and vulnerable populations [2]. These are achievable by strengthening the health system for safety and enhanced potential of lives and livelihood through conscientious efforts, restorative and sustainable [3] financing and technical capacity.

It is pertinent that governments and other contributing agents provide the general public with expansive information and communication needs for the treatment and control of the veritable symptoms regarding SARS-CoV-2 or COVID-19 [4, 5] in order that global citizens taper or curb the never-ending prevalent scepticism. Procrastinations in minimizing restrictions have been associated with incremental vaccinations, dissemination, transmission and upsurge of the new Delta variant that is perspicuously overwhelming the health system and becoming a global scourge.

Outlook

Even with this dimension, vaccination intensity comparatively declined spatiotemporally, partially due to the exclusion of those below the fourth decade of life as beneficiaries/recipients of the AstraZeneca version and the preponderant reluctance to be vaccinated. It is, however, predicted that the Delta variant has disseminated involving circa 50,000 new infections daily within a short period in the instant year. The critical aspect may be effectively and efficiently circumvented by curbing the untoward impact of the variants via vaccine combination or multifactorial applications, mask adornment, social distancing and invariable lockdown as in the herd community strategy [6].

In Nigeria, the presence of the Delta variant called lineage B.1.617.2 of the COVID-19 or SARS-CoV-2 virus was revealed [7]. It is suggested that the Delta variant is more adverse than the Alpha variant with double the latitude for infection, hyperinfection, delayed convalescence, increased dissemination, debilitating outcome or resultant impact and concomitant hospitalization. It becomes imperative to inter alia adhere to appropriate guidelines in the applications of nose, mouth and face masks in crowded ambients, handwashing with soaps and running water as well as hand sanitizer usage. It has become advisable that travelers from outside Nigeria undergo the mandatory seven-day isolation period with concomitant testing prior to physical communication with the general public. Residents are advised to seek medical care/healthcare services, and avail themselves of free-testing opportunities where there are certainties of resultant morbidity and mortality, as measures to ameliorate or curb the dissemination and preponderance of SARS-COV-2 or COVID-19 in the environs. The National Primary Health Care Development Agency, NPHCDA states that Nigeria will be the beneficiary/recipient of 41,282,770 doses of Oxford/AstraZeneca, Pfizer-Bio-N-Tech and Johnson & Johnson COVID-19 or SARS-CoV-2 vaccines within July and September this year.

Ignoring preventive strategies against the COVID-19 places everybody at risk for infection. Shortly, following the confirmation of the Delta variant in Nigeria, cases have consistently become elevated in the country infecting unprotected and vulnerable persons with accelerated pressure on the deficient provision of oxygen, therapeutics, healthcare systems and services, personnel protective equipment, exasperated health workers, low income, and other shortages paving the trajectory for deteriorating public health emergency with the resultant threat to human life, livelihood, healthy economy, restoration and sustainability [7].

Etiopathogenesis and Vaccinations

A study [8] depicts a key alteration that placed the Greater Horse-shoe coronavirus bat, *Rhinolophus ferrumequinum* on the trajectory for

infection with a pathogenic impact on humans from animals. The mutation ostensibly aids the virus per the spike protein, and adhere to the human replica of a host protein, hACE2, thus creating the latitude for the virus to penetrate and infect cells. The mutated virus has a greater potential lock and adhere to human cells than other coronaviruses deficient of the alterations; and also has greater replicative capacity in experimental human lung cells than erstwhile versions of the coronavirus. It is, however, suggested that this does not necessarily connote the sole mutation, and that the strategy employed in the research is not conventionally applicable in virology with the tendency not to have detected other pertinent or appreciable mutations. However, the COVID-19 or SARS-CoV-2 culprit is the substitution of the amino acid, threonine that is ubiquitous in animal viruses by its counterpart, alanine. The mutation, T372A may have been due to the deletion of certain sugars which coat the spike protein. These sugars probably impede pathogenicity, thus their eradication provides the latitude for hACE2 to permeate cells. An excess of 182,000 SARS-COV-2 genomes screened for selective sweep signatures depicted an adaptive alteration with the spike protein receptor-binding domain. The alteration was predicted, and the laboratory-setting correlated to enhanced affinity to hACE2 with resultant viral replication in relation to conventional ancestral variant. The culminating mutation ostensibly contributes to the emergence of SARS-COV-2 from animal reservoirs with potentiated transmission to humans [8].

Thus, mRNA vaccines coated with soft fatty acids utilise genetic material in addition to diverse inactive excipients to construct an aspect of the coronavirus spike protein, and directs the immune system for the formation of specific antibodies [9]. The soft lipids permit the mRNA to gain entry into cells for cytoplasmic absorption and initiation of the spike protein synthesis. Predictably, vaccination activates T cells which potentiate the immune system response to future coronavirus exposure [9]. The mRNA induces SARS-COV-2 or COVID-19 antigen synthesis that triggers the antibody response of the vaccinated individual via the formation of neutralizing antibodies.

The COVID-19 or SARS-CoV-2 variants

Several strains of the coronavirus have been identified in recent outbreaks. The Delta variant depicts that vaccines are effective in the dilemma of the COVID-19 or SARS-CoV-2 pandemic. In ambients with infinitesimal COVID-19 or SARS-CoV-2 vaccine coverage there have been an upsurge of morbidity, mortality and expansively overwhelmed healthcare infrastructure including gross economic downturn [2]. These correlate with the Delta variant (B.1.617.2) dominance [10]. This strand presents greater transmissibility, resilience, and suppresses vaccine efficacy. The dissemination of variants, such as B.1.1.7 and B.1.351 has prompted extensive effort to prevent and control the SARS-COV-2 or COVID-19 pandemic through vaccination through disparate modalities for emerging variants [11]. A procrastinated vaccine rollout has resulted in adverse repercussions in certain countries. However, after vaccinating the risk groups, Israel altered her itinerary in this COVID-19 or SARS-CoV-2 pandemic era in its policy of "soft suppression" that entails living with the virus by ameliorating expansive restrictions and a fourth national lockdown which could perspicuously impede the economic potential of the country [12]. The opportunity cost may be the ravaging tendency of the incipient variants.

The Lambda variant of COVID-19 or SARS-CoV-2 was declared by

WHO as a "Variant of Concern", VOC. The Lambda strand or variant C.37 was initially detected in Peru [13], and classified as a "Variant of Interest" by WHO. A VOI is defined as a variant that is commonly established in diverse communities with mutations which give rise to certain alterations within the virus exhibiting atypical symptoms. A VOI exhibits mutations which are predicted or realised to impact factors, such as confounding of diagnostic tests, transmissibility, disease severity, and propensity to elude immunity from erstwhile infections or vaccines. Variant of Concern merely specifically relates that the variant has the attribute of elevated rate of transmission. Although, the Lambda variant has been established in several countries, excepting those in Africa and Asia, it is not yet classifiable as a VOC because of deficient evidence in its transmission potential. It is, however, designated as a "Variant Under Investigation" due to its "international expansion and several notable mutations".

In retrospect, the Lambda variant is a coronavirus strain that has recently been detected in about 30 countries. It is associated with extensive community transmission rates in numerous countries, with spatiotemporally increasing prevalence and increased incidence of COVID-19 or SARS-CoV-2. The unusual combination of mutations due to the Lambda variant may constitute its greater transmissibility. There is a paucity of data or evidence to suggest that these combinatorial mutations depict the Lambda variant as being more adverse than the originating coronavirus. The Pfizer and Moderna vaccines are protective against the Lambda variant, and there is no evidence to indicate that it has greater pathogenic virulence than the Delta variant. Thus, it remains of interest and not of concern [13]. The Delta strain presents not merely greater transmissibility but its infection rate deleteriously impacts on erstwhile COVID-19 or SARS-CoV-2 subjects. Currently, vaccination becomes imperative as every instant and extant COVID-19 or SARS-CoV-2 is a preventable mortality. A recent Kappa variant that is also more benign than its Delta counterpart has emerged in Rajasthan [14].

Perspectives

A vast majority of viral mutations may not impact on the attributes of the pathogenicity of the viruses in their dissemination, transmission and severity depending on the locale of alterations in their genetic materials. The extant COVID-19 or SARS-CoV-2 vaccines on implementation are predicted to provide certain or absolute protection against novel variants due to the potential of the vaccines to elicit expansive immune response in concert with varied cells and antibodies. In the modicum of events that alterations or mutations attenuate or debilitate the effectivity of a vaccine, the components of the vaccine need to be fortified or modified for efficacy, effectivity and efficiency. Implementation of public health policies must not detract concerns of SARS-CoV-2 or COVID-19 variants as to diminish vaccine efficacy per VOCs culminating in excoriated vaccination rates of extant SARS-CoV-2 or COVID-19 [15], and as the spike protein portend constraints for monoclonal antibody treatment and pose bottlenecks to the efficacy of extant vaccines [16]

Discussion

Besides other anomalies and untoward events herein brought to the fore, the COVID-19 or SARS-CoV-2 pandemic has culminated in adverse misinformation [17]. There are spatiotemporal variations and changes regarding viruses and other microorganisms [18] which are characteristic

of COVID-19 or SARS-CoV-2. The concomitant changes may not translate into pathogenic mutants which adversely impact on the attributes of the viruses in certain instances per dissemination, transmission, associated pathogenicity, reliability of vaccines, diagnostic and other social, environmental and public health strategies.

It is mandatory to establish, improve and undergird extant social and public health programmes as well as infection prevention and control strategies for effective and efficient diminution of COVID-19 or SARS-CoV-2 virulence, prevalence, incidence, hospitalisation, morbidity and mortality. These are realisable by undergirding surveillance, sequencing capacities, and applications of systematic modalities to predict the magnitude of dissemination and transmission of SARS-CoV-2 or COVID-19 variants by thinking locally and acting globally in order to preclude the incident risk of the pandemic for restoration and sustainability [3].

There are tendencies for several countries to lift, ease, continue or augment restrictions as they battle the COVID-19 or SARS-CoV-2 pandemic despite the accelerated dissemination of the Delta variant. Globally, there are perspicuous economic prospects [2, 19] with availability of vaccines combined with easing or discontinuance of restrictions, especially in the education, tourism, hospitality, social, recreational, industrial, manufacturing and production sectors. These considerations must, therefore, not merely relate to the public health perspective despite the superimposed economic preponderance.

The successful achievement of the vaccination programme will negatively correlate the number of cases, hospitalisations and mortality, even with disproportionate number of individuals who elect to decline vaccination or further jabs. As there are those in the industrialized countries who are ostensibly presented with the choice of being vaccinated, this choice may not be available to most vulnerable populations in less developing countries. These susceptible individuals are at risk of being overwhelmed by novel variants and decreased restrictions.

Conclusion

Global production, distribution and consumption of vaccines and booster doses to curb the scourge of the SARS-CoV-2 or COVID-19 pandemic with its perspicuous incessant and proliferative variants are grossly inequitable within certain countries and vulnerable populations. It is desirable to have access to increased volume of vaccines as they portend prolonged immunity against untoward and life-threatening COVID-19 SARS-CoV-2 virulence rather than prioritizing mere restrictions and lockdowns combined with other COVID-19 or SARS-CoV-2 protocols. The attendant consequences of the opportunity cost of not presenting an encompassing multifactorial strategy of curbing the cascade of variants via decisions on access to vaccines, booster doses and restrictions must be obviated.

References

- [1] Choudhary OP, Priyanka, Singh I and Rodriguez-Morales AJ. Second wave of COVID-19 in India: Dissection of the causes and lessons learnt. *Travel Med Infect Dis.* 2021; 43: 102126. DOI: 10.1016/j.tmaid.2021.102126.
- [2] Chukwuma Sr C. Re-thinking poverty and health issues in the environment of the COVID-19 pandemic. *Int J Adv Res Rev, IJARR.* 2021; 6(1): 26-29.
- [3] Chukwuma Sr C. Ecological analysis of the COVID-19 pandemic for restoration and sustainability. *Research and Advances in Pharmacy and Life Sciences.* 2021; Vol 3, Issue 1.
- [4] Priyanka, Choudhary OP, Singh I, Patra G. Aerosol transmission of SARS-CoV-2: The unresolved paradox. *Travel Med Infect Dis.* 2020; 37: 101869. DOI: 10.1016/j.tmaid.2020.101869.
- [5] Chukwuma Sr C. Information and communication needs in the administration of ibuprofen in the treatment of symptoms in COVID-19 patients. *Journal of Medical Research, JMR.* 2020; 6(4): 128-130. <https://doi.org/10.31254/jmr.2020.6405>.
- [6] It's time the government levelled with us about Covid and the science behind 'freedom day'. www.independent.co.uk/voices/letters/covid-vaccination-freedom-day-letters-b1882087.html.
- [7] ALERT: Deadly COVID-19 strain discovered in Oyo. <https://politicsnigeria.com/alert-deadly-covid-19-strain-discovered-in-oyo/>.
- [8] Kang L, He G, Sharp AK, Wang X et al. A selective sweep in the Spike gene has driven SARS-COV-2 human adaptation. <https://doi.org/10.1016/j.cell.2021.07.007>.
- [9] Conti P, Caraffa A, Gallenga CE, Kritas SK, Frydas I, Younes A et al. The British variant of the new coronavirus-19 (SARS-COV-2) should not create a vaccine problem. *J Biol Regul Homeost Agents.* 2021; 35(1): 1-4. doi: 10.23812/21-3-E.
- [10] COVID-19 delta variant spreads in countries with low vaccinations. <https://qz.com/india/2032726/covid-19-delta-variant-spreads-in-countries-with-low-vaccinations/>.
- [11] Bian L, Gao F, Zhang J, He Q, Mao Q, Xu M, Liang Z et al. Effects of SARS-COV-2 variants on vaccine efficacy and response strategies. *Expert Rev Vaccines.* 2021; 1-9. <https://doi.org/10.1080/14760584.2021.1903879>.
- [12] Israel changes strategy as delta variant makes 90 of the cases. <https://www.euractiv.com/section/coronavirus/news/israel-changes-strategy-as-delta-variant-makes-90-of-the-cases/>.
- [13] Risk Assessments for SARS-COV-2 variant Delta 8 July, 2021. Ref: PHE publications gateway number: GOV-8894.
- [14] 11 cases of new kappa variant of COVID-19 detected in Rajasthan. <https://www.newindianexpress.com/nation/2021/jul/14/11-cases-of-new-kappa-variant-of-covid-19-detected-in-rajasthan-2329706.html>.
- [15] Noh JY, Jeong HW, Shin E-O. SARS-COV-2 mutations, vaccines and immunity: implications of variants of concern. *Sig Trans Targ Ther.* 2021; 6, Article No.: 203. <https://dou.org/10.1038/s41392-021-00623-2>.
- [16] Wang P, Nair MS, Liu L et al. Antibody resistance of SARS-COV-2 variants B:1.351 and B.1.1.7. *Nature.* 2021; 593: 130-135. <https://doi.org/10.1038/s41586-021-03398-2>.
- [17] Misleading. <https://www.logically.ai/factchecks/library/c3b1eec1>.
- [18] Chukwuma Sr C. Bioinformatics-base and determinants in the spatiotemporal variations of emerging and reemerging infectious diseases. *Journal of Ancient Diseases & Preventive Remedies.* 2018; 06(02). DOI: 10.4172/2329.8731.1000182.
- [19] Chukwuma Sr C. Strategy on poverty exacerbated by the cCOVID-19 pandemic. *IjCSRSD.* 2021; 1(2): 105-107.