

CO-ORGANIZED EVENT

International Conference on **Chronic Diseases**

&
6th International Conference on **Microbial Physiology and Genomics**

August 31-September 01, 2017 Brussels, Belgium

Detection of JC Polyomavirus Tumor-Antigen In Gastric Carcinoma, a report from IRAN

Farnaz Zahedi Avval^{1,2}, Samira Izi¹, Farzad Rahmani¹, Nema Mohammadian Roshan³, Atefeh Yari⁴, Masoud Youssefi^{4,5}

¹ Department of Clinical Biochemistry, School of Medicine, Mashhad University of Medical Sciences, Mashhad, IRAN

² Metabolic Syndrome Research Center, Nutrition and Biochemistry division, Mashhad University of Medical Sciences, Mashhad, IRAN

³ Department of Pathology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, IRAN

⁴ Department of Microbiology and Virology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, IRAN

⁵ Antimicrobial Resistance Research Center, Mashhad University of Medical Sciences, Mashhad, IRAN

Background: Although the global incidence of gastric cancer has been declined dramatically over the recent few decades, it is still a common cancer in different parts of the world. It has been scientifically suggested that some oncogenic viruses like JCV might play a role in Gastric carcinogenesis. The viral transforming protein, T-antigen (T-Ag) has the ability to bind and inactivate tumor suppressor proteins including p53 and pRb that might lead to malignant outcome, though its role might vary in different geographic parts due to viral distribution and other habits resulting in GI cancer.

Objective: The aim of present study was to investigate the presence of JCV T-Ag sequence and its expression in cancerous and non-cancerous adjacent gastric tissues in Iranian patients.

Methods: Thirty one sample pairs of formalin fixed paraffin embedded (FFPE) tissue specimens of gastric cancer and adjacent non-cancerous tissues (ANCT) were investigated on the basis of Real-time polymerase chain reaction. Samples were subjected for the immunohistochemistry examination using an anti-T-Ag monoclonal antibody.

Results: Real time experience followed by sequencing revealed JCV sequences in 17 (54.84%) of gastric cancer tissues and in 10 (32.25%) of non-cancerous gastric mucosa (OR=1.7). Immunohistochemical study also showed the presence of T-Ag in nuclear compartment.

Conclusions: These data indicate, for the first time, presence of JC virus in gastric carcinoma samples in our socioeconomic region. These findings provide a summative, supportive data for a possible role of JCV T-Ag in carcinogenesis of gastric malignancy.

ZahediAF@mums.ac.ir

Notes: